

Almost positive links have negative signature

Józef H. Przytycki and Kouki Taniyama

Draft 1990-1991; Corrections July-September 1999; Completion May-July 2008,
February-April 2009

Abstract.

We analyze properties of links which have diagrams with a small number of negative crossings. We show that if a nontrivial link has a diagram with all crossings positive except possibly one, then the signature of the link is negative. If a link diagram has two negative crossings, we show that the signature of the link is nonpositive with the exception of the left-handed Hopf link (with possible trivial components). We also characterize those links which have signature zero and diagrams with two negative crossings. In particular, we show that if a nontrivial knot has a diagram with two negative crossings then the signature of the knot is negative, unless the knot is a twist knot with negative clasp. We completely determine all trivial link diagrams with two or fewer negative crossings. For a knot diagram with three negative crossings, the signature of the knot is nonpositive except the left-handed trefoil knot. These results generalize those of L. Rudolph, T. Cochran, E. Gompf, P. Traczyk, and J. H. Przytycki, solve Conjecture 5 of [P-2], and give a partial answer to Problem 2.8 of [Co-G] about knots dominating the trefoil knot or the trivial knot. We also describe all unknotting number one positive knots.

1. INTRODUCTION

This paper is a sequel to [P-2] and [T-1, T-2] and uses ideas from these papers. K. Murasugi, following C. A. Giller [Gi] showed that nontrivial positive alternating links have negative signature. In 1982 L. Rudolph showed that nontrivial positive braids have negative signature (in our orientation convention), [R-1]. These results are generalized in [Co-G, Tr, P-2] to the theorem that nontrivial positive links have negative signature. This theorem was first proven (for knots) by T. Cochran and R. Gompf in the summer of 1985 (at MSRI) but the paper [Co-G] was written only two years later. The theorem was independently proven by P. Traczyk in summer of 1987 [Tr], and by J. H. Przytycki in September of 1987 [P-2]. In this paper¹ we generalize the theorem to show that nontrivial almost positive links and 2-almost positive links (except twist knots, the left-handed Hopf link, and the connected or disjoint sum of the left-handed Hopf link, and a $(2, 2k)$ -torus link) have negative signature.

¹Collaboration on this paper began when Przytycki noticed that the affirmative answer to a question he asked in 1987 [P-2] followed from Taniyama's master thesis (see [T-1, T-2]). The question was whether nontrivial almost positive link has negative signature. The first draft of the paper was written in 1990-1991.

To describe precisely the content of our paper we start with a few preliminary definitions.

We consider oriented links in $\mathbb{S}^3$ and oriented link diagrams on $\mathbb{S}^2$ up to ambient isotopy of $\mathbb{S}^3$ and $\mathbb{S}^2$ respectively. A link diagram is *m-almost positive* if all but m of its crossings are positive ($\times$).

A 0-almost positive diagram is called positive and a 1-almost positive diagram is called almost positive. A link is *m-almost positive* if it has an *m-almost positive* diagram. In particular a 0-almost positive link is called positive and a 1-almost positive link is called almost positive. For links L_1 and L_2 , we define the relation $\geq$ as follows: L_1 is greater than or equal to L_2 , denoted $L_1 \geq L_2$, if L_2 can be obtained from L_1 by changing some positive crossings to negative crossings. This relation is weaker than the relation in [Co-G] and different from, but related to, the relation of [T-1]. We denote by $\preceq$ the relation defined on knots in [Co-G].² Here we only need the fact that if $K_1 \geq K_2$ then $K_1 \succeq K_2$. It was observed by Giller [Gi] and utilized by Murasugi, Cochran and Lickorish [Co-Li] that if links L_1 and L_2 are oriented links which differ only by one crossing at which L_1 is positive and L_2 is negative then,

$$\sigma(L_2) - 2 \leq \sigma(L_1) \leq \sigma(L_2)$$

where $\sigma(L)$ denotes the signature of L . Thus for links L_1 and L_2 with $L_1 \geq L_2$ we have the inequality $\sigma(L_1) \leq \sigma(L_2)$. We call this the Giller inequality in our paper³. This inequality is a basic tool used in the paper. In Section 2 we work with positive links. There we will prove the following result:

Theorem 1.1. *Let K be a nontrivial positive knot. Then $K \geq (2,5)$ -torus knot or K is a connected sum of some pretzel knots of type $L(p_1, p_2, p_3)$, where p_1, p_2 and p_3 are positive odd numbers; see Fig. 1.1.*




FIGURE 1.1.

Corollary 1.2. *Let K be a nontrivial positive knot. Then $K \succeq (2,5)$ -torus knot or K is a connected sum of some pretzel knots of type $L(p_1, p_2, p_3)$, where p_1, p_2 and p_3 are positive odd numbers.*

² According to [Co-G], $K_1 \succeq K_2$ if K_1 is concordant to K_2 inside of a 4-manifold with positive definite intersection form whose concordance annulus is homologically trivial. See Definition 2.1 in [Co-G].

³Giller was motivated by J. H. Conway [Con], and used the inequality to show that the signature of knots is a skein equivalence invariant. Conway, in turn, was influenced by Murasugi's papers on signature [M-1, M-2].

Corollary 1.3. *Let K be a nontrivial positive knot. Then either the signature $\sigma(K) \leq -4$, or K is a pretzel knot $L(p_1, p_2, p_3)$, where p_1, p_2 and p_3 are positive odd numbers. We have $\sigma(L(p_1, p_2, p_3)) = -2$.*

In Section 3 we analyze almost positive links. We prove there in particular:

Theorem 1.4. *Let L be a nontrivial almost positive link. Then $L \geq$ right-handed trefoil knot (plus trivial components), or $L \geq$ right-handed Hopf link (plus trivial components).*

Theorem 1.5. *Let $\tilde{L}$ be an almost positive diagram representing a trivial link. Then $\tilde{L}$ can be reduced to one of the diagrams of Fig. 1.2 by first Reidemeister moves (reducing the number of crossings), and deleting trivial circles.*




FIGURE 1.2. Almost positive diagrams representing a trivial link

Corollary 1.6. *Let K be a nontrivial almost positive knot. Then $K \succeq$ right-handed trefoil knot.*

Corollary 1.7. *Nontrivial almost positive links have negative signature.*

In Section 4 we analyze 2-almost positive links. In particular, we prove:

Theorem 1.8. *Let $\tilde{L}$ be a nontrivial 2-almost positive link. Then*

- (1) $L \geq$ right-handed trefoil knot (plus trivial components), or
- (2) $L \geq 6_2$ (Fig. 1.4 (a)) (plus trivial components), or
- (3) $L \geq$ right-handed Hopf link (plus trivial components), or
- (4) $L \geq$ disjoint or connected sum of right-handed trefoil knot and left handed Hopf link (plus trivial components), or
- (5) $L \geq$ Whitehead link (Fig. 1.4 (b)) (plus trivial components), or
- (6) $L \geq$ disjoint and/or connected sum of two right-handed Hopf links and a left-handed Hopf link (plus trivial components), or
- (7) $L \geq$ disjoint or connected sum of (2,4)-torus link (Fig. 1.4 (c)) and a left-handed Hopf link (plus trivial components), or
- (8) $L \geq$ the link of Fig. 1.4 (d) (plus trivial components), or
- (9) $\tilde{L}$ is a twist knot with negative clasp (Fig. 1.3) (plus trivial components), or
- (10) L is a disjoint or connected sum of left-handed Hopf link and (2,n)-torus link with anti-parallel orientation of components (Fig. 1.5) (plus trivial components), or
- (11) L is a left-handed Hopf link (plus trivial components).




FIGURE 1.3. Twist knots with negative clasp





FIGURE 1.4.

FIGURE 1.5. $(2, n)$ -torus link with anti-parallel orientation of components

Theorem 1.9. *Let $\tilde{L}$ be a 2-almost positive diagram (drawn on $S^2 = R^2 \cup \infty$) representing a trivial link. Then $\tilde{L}$ can be obtained from the diagrams in Fig. 1.2 and Fig. 1.6, or their π -rotation along y axis, by performing some combination of diagram disjoint sum operation, diagram connected sum operation, and first and second Reidemeister moves which increase the number of crossings.*

Corollary 1.10. *A nontrivial 2-almost positive link L has nonnegative signature if and only if*

- (1) *L is a twist knot with negative clasp (with, possibly, trivial components), in which case $\sigma(L) = 0$, or*
- (2) *L is a left-handed Hopf link (with, possibly, trivial components), in which case $\sigma(L) = 1$, or*
- (3) *L is a disjoint or connected sum of left-handed Hopf link and $(2, n)$ -torus link with anti-parallel orientation of components (Fig. 1.5) (with, possibly, trivial components), in which case $\sigma(L) = 0$.*

In Section 5 we consider 3-almost positive knots and prove:

Theorem 1.11. *Let K be a 3-almost positive knot. Then either $K \geq$ trivial knot or K is the left-handed trefoil knot (plus positive knots as connected summands).*

Corollary 1.12. *Let K be a 3-almost positive knot, then*




FIGURE 1.6.

- (a) either $\sigma(K) \leq 0$ or K is the left-handed trefoil knot.
- (b) either $K \succeq$ trivial knot or K is the left-handed trefoil knot (plus positive knots as connected summands).

In Section 6 we use our previous results to characterize the unknotting number one positive links, 2-almost positive amphicheiral links, and 2-almost positive slice links. We also analyze homology 3-spheres not bounding a (rational) homology 4-ball, the Tristram-Levine signatures and the Jones polynomial. In particular, we prove:

- Theorem 1.13.**
- (a) If a positive knot has unknotting number one then it is a positive twist knot.
 - (b) If K is a 2-almost positive knot different from a twist knot with a negative clasp then $K(1/n)$ (i.e. $1/n$ surgery on K , $n > 0$) is a homology 3-sphere that does not bound a compact, smooth homology 4-ball. Furthermore, $K(1/n)$ has a nontrivial Floer homology.
 - (c) If K is a non-trivial 2-almost positive knot different from the stevedore's knot then K is not a slice knot.

- (d) If K is a non-trivial 2-almost positive knot different from the figure eight knot then K is not amphicheiral.

2. POSITIVE LINKS

It has been proved in [Co-G] and [P-1] that if K is a nontrivial positive knot then $K \geq$ right-handed trefoil knot. We generalize this result in this and the next sections.

2.1. Preliminary terminology. If we ignore over/under crossing information of a link diagram $\tilde{L}$, then we call it the underlying *projection* (or *universe*; [Ka-1]) of $\tilde{L}$, and denote it by $\hat{L}$. In general $\tilde{X}$ denotes some part of a diagram and $\hat{X}$ denotes its underlying projection.

A tangle is composed of properly embedded (oriented) system of arcs in a 3-ball $\mathbb{B}^3$. We consider tangles up to ambient isotopy of $\mathbb{B}^3$ which is fixed on $\partial\mathbb{B}^3$. For oriented tangles T_1 and T_2 , we define the relation $\geq$ where $T_1 \geq T_2$ if T_2 can be obtained from T_1 by changing some positive crossings to negative crossings. We consider tangle diagrams $\tilde{T}$ and its underlying projection $\hat{T}$ on the unit disk $\mathbb{B}^2$ up to ambient isotopy which is fixed on $\partial\mathbb{B}^2$. Note that no tangles, tangle diagrams and tangle projections in this paper contain closed components unless otherwise stated. For a link projection $\hat{L}$ (resp. tangle projection $\hat{T}$) we denote by $\text{LINK}(\hat{L})$ (resp. $\text{TANGLE}(\hat{T})$) the set of all links (resp. tangles) that has a link diagram (resp. tangle diagram) whose underlying projection is $\hat{L}$ (resp. $\hat{T}$). For two link (resp. tangle) projections $\hat{L}_1$ and $\hat{L}_2$ (resp. $\hat{T}_1$ and $\hat{T}_2$), we define the relation $\hat{L}_1 \geq \hat{L}_2$ (resp. $\hat{T}_1 \geq \hat{T}_2$) if $\text{LINK}(\hat{L}_1) \supset \text{LINK}(\hat{L}_2)$ (resp. $\text{TANGLE}(\hat{T}_1) \supset \text{TANGLE}(\hat{T}_2)$). Then we say that $\hat{L}_2$ (resp. $\hat{T}_2$) is a *minor* of $\hat{L}_1$ (resp. $\hat{T}_1$).

An oriented tangle diagram is *m-almost positive* if all its crossings, but exactly m , are positive.

A crossing P of $\tilde{L}$ (resp.⁴ $\tilde{L}, \tilde{T}, \hat{T}$) is called *nugatory* if $\hat{L} - P$ (resp. $(\partial\mathbb{B}^2 \cup \hat{T}) - P$) has more components than $\hat{L}$ (resp. $\partial\mathbb{B}^2 \cup \hat{T}$). A link (resp. tangle) diagram without nugatory crossings is called *reduced*.

A link (resp. tangle) diagram is called *R2-reduced* if it is reduced and it does not contain a 2-gon; as illustrated in Fig. 2.1




FIGURE 2.1.

We say that a link (resp. tangle) diagram is *prime* if it is connected and any simple closed curve in $\mathbb{S}^2$ (resp. $\mathbb{B}^2$) meeting it transversally in two points bounds

⁴ Most of the definition we give below apply to the case of a link diagram $\tilde{L}$, its projection $\hat{L}$, and a tangle diagram $\tilde{T}$ and its projection $\hat{T}$. To omit cumbersome notation/repetitions we will list only one or two of them in definitions, unless description differs, in which case we list the difference.

a trivial disk pair. We remark that prime link projections are reduced except a projection of a trivial knot with one crossing.

Let $\hat{L}$ be a link projection that is not prime. Let γ be a simple closed curve on $\mathbb{S}^2$ meeting it transversally in two points, say P and Q , that does not bound trivial disk pair. Let D_1 and D_2 be disks in $\mathbb{S}^2$ bounded by γ . Let α be a simple arc in γ joining P and Q . Let $\hat{L}_1 = (\hat{L} \cap D_1) \cup \alpha$ and $\hat{L}_2 = (\hat{L} \cap D_2) \cup \alpha$. Then we say that $\hat{L}$ decomposes to $\hat{L}_1$ and $\hat{L}_2$. We also say that $\hat{L}$ is a *connected sum* of $\hat{L}_1$ and $\hat{L}_2$. See Fig. 2.2.




FIGURE 2.2.

A *self-crossing* of a link (resp. tangle) diagram is a crossing involving a single component. A *mixed crossing* is a crossing between two different components.

We say that two mixed crossings on a component are *successive* on the component if there is an arc in the component connecting the crossings without mixed crossings on the arc.

A self-crossing P of an arc component $\tilde{t}$ of a tangle diagram $\tilde{T}$ divides $\tilde{t}$ into three immersed arcs. Let $r(\tilde{t}, P)$ be the diagram obtained from $\tilde{t}$ by removing the middle immersed arc.⁵ Let $s(\tilde{t}, P)$ be the middle part so that $\tilde{t} = r(\tilde{t}, P) \cup s(\tilde{t}, P)$. Let $r(\tilde{T}, P)$ be the diagram obtained from $\tilde{T}$ by replacing $\tilde{t}$ by $r(\tilde{t}, P)$. Similarly, a self-crossing P of a component $\tilde{\ell}$ of a link diagram $\tilde{L}$ divides $\tilde{\ell}$ into two immersed arcs. We denote one of them by $r(\tilde{\ell}, P)$ and the other by $s(\tilde{\ell}, P)$. In a similar manner we define $r(\tilde{L}, P)$ and $s(\tilde{L}, P)$.

A component $\tilde{\ell}$ (resp. $\tilde{t}$) of a link (resp. tangle) diagram is said to be *almost trivial* if $\tilde{\ell} - P$ (resp. $\tilde{t} - P$) is not connected for any self-crossing P of $\tilde{\ell}$ (resp. $\tilde{t}$).⁶

For a component $\tilde{t}$ of a tangle diagram $\tilde{T}$ and a component (arc) α of $\partial\mathbb{B}^2 - \partial\tilde{t}$, we say that a point P on $\tilde{t}$ is *outermost with respect to α* if α and P belong to the same closure of a component of $\mathbb{B}^2 - \tilde{t}$. When α contains the point $(1, 0)$ (resp. $(-1, 0)$) we say that $\tilde{t}$ is *rightmost* (resp. *leftmost*).

For a point A on $\tilde{t}$ and a component α of $\partial\mathbb{B}^2 - \partial\tilde{t}$, we define the *depth* of A on $\tilde{t}$, $d(A) = d(A, \alpha)$, to be the minimal number of the transverse intersection points of $\tilde{t}$ and an arc joining α and A (avoiding crossing points) which is in general position with respect to $\tilde{t}$. In particular $d(A) = 0$ if A is outermost. When α contains the point $(1, 0)$ (resp. $(-1, 0)$) we denote $d(A, \alpha) = d(A, (1, 0))$ (resp. $d(A, \alpha) = d(A, (-1, 0))$) and call it the *right depth* (resp. *left depth*) of A on $\tilde{t}$.

For an oriented arc component $\hat{t}$ of a tangle diagram $\hat{T}$, we introduce inductively the following terminology. Let Γ_0 be the set of all self-crossings of $\hat{t}$. Set $\hat{t}_0 = \hat{t}$.

⁵That is, we smooth the crossing P and delete the closed component.

⁶That is, a knot $\tilde{\ell}$ (resp. an arc $\tilde{t}$) has only nugatory crossings.

We trace $\hat{t}_0$ along its orientation starting from its first end point. Let P_1 be the first self-crossing of $\hat{t}_0$ we encounter. Let $\hat{t}_1 = r(\hat{t}_0, P_1)$. Then we trace $\hat{t}_1$ starting from P_1 along its orientation. Let P_2 be the first self-crossing of $\hat{t}_1$ we encounter. Let $\hat{t}_2 = r(\hat{t}_1, P_2)$. We continue this and finally have $\hat{t}_{n_1}$ that has no self-crossings. We set $\hat{t}' = \hat{t}_{n_1}$ and call it the *spine* of $\hat{t}$.

Let Γ_1 be $\Gamma_0 - \hat{t}'$. Namely Γ_1 is the set of self-crossings of $\hat{t}$ that are not on $\hat{t}'$.

Set $\hat{t}'_0 = \hat{t}'$ again. We trace again $\hat{t}'_0$ along its orientation starting from its first end point. Let P'_1 be the first self-crossing in Γ_1 we encounter. Let $\hat{t}'_1 = r(\hat{t}'_0, P'_1)$. Then we trace $\hat{t}'_1$ starting from P'_1 along its orientation. Let P'_2 be the first self-crossing of $\hat{t}'_1$ in Γ_1 we encounter. Let $\hat{t}'_2 = r(\hat{t}'_1, P'_2)$. We continue this and finally have $\hat{t}'_{n_2}$ whose all self-crossings are on $\hat{t}'$. We set $\hat{t}'' = \hat{t}'_{n_2}$ and call it the *second spine* of $\hat{t}$.

We define inductively the n -th spine $\hat{t}^{(n)}$ in a similar manner. In particular we define $\Gamma_n = \Gamma_{n-1} - \hat{t}^{(n)}$. Of course $\hat{t}^{(n-1)} \subset \hat{t}^{(n)}$.

See for example Fig. 2.3.




FIGURE 2.3.

For a point A in an almost trivial arc component $\hat{t}$, we define the *multiplicity* of A on $\hat{t}$, $m(A) = m(A, \hat{t})$ to be the minimal number n such that A is contained in $\hat{t}^{(n+1)}$. A self-crossing P of $\hat{t}$ is called the *root* of A if P is on $\hat{t}^{(m(A))}$ and A is on $s(\hat{t}, P)$. Note that every A with $m(A) \geq 1$ has a unique root. For points A and B in $\hat{t}$, we say that A and B are *related on $\hat{t}$* if $m(A) = m(B)$ and A and B belong to the same component of $\hat{t} - (\Gamma_0 - \Gamma_{m(A)})$. Note that if $m(A) = m(B) \geq 1$ then they have the common root P . See for example Fig. 2.4.




FIGURE 2.4.

We say that a link diagram $\tilde{L}_1$ (resp. $\tilde{T}_1$) is an *R1 augmentation* of a link diagram $\tilde{L}_0$ (resp. $\tilde{T}_0$) if $\hat{L}_1$ (resp. $\hat{T}_1$) can be obtained from $\hat{L}_0$ (resp. $\hat{T}_0$) by a finite sequence of the first Reidemeister moves increasing the number of crossings as illustrated in Fig. 2.5.




FIGURE 2.5.

If $\tilde{L}_1$ is an R1 augmentation of $\tilde{L}_0$ then $\hat{L}_0$ is a subspace of $\hat{L}_1$ and we say that $\hat{L}_1 - \hat{L}_0$ is an *R1 residual* of $\hat{L}_1$ with respect to $\hat{L}_0$, and $\hat{L}_0$ is a *core* of $\hat{L}_1$.

Throughout the paper we use the notation introduced in Fig. 2.6 for a 2-string tangle diagram (resp. projection). Namely the endpoints of the strings are denoted by A_0, A_∞, B_0 and B_∞ , and the arc component A_0A_∞ is denoted by $\tilde{a}$ (resp $\hat{a}$), and the arc component B_0B_∞ is denoted by $\tilde{b}$ (resp $\hat{b}$). Recall that $\tilde{x}$ denotes a diagram and $\hat{x}$ denotes a projection. Let $\tilde{a}'$ be the spine of $\tilde{a}$ and $\tilde{b}'$ the spine of $\tilde{b}$. We call $\tilde{T}' = \tilde{a}' \cup \tilde{b}'$ the *spine* of $\tilde{T} = \tilde{a} \cup \tilde{b}$.




FIGURE 2.6.

We say that the tangle diagram (resp. projection) of Fig. 2.6 (a) has a *vertical connection* and of Fig. 2.6 (b) has an *X-connection*. We also sometimes use the tangle diagram (resp. projection) of Fig. 2.6 (c) and say it has a *horizontal connection*. The string is always oriented from A_0 to A_∞ and from B_0 to B_∞ unless otherwise stated.

We denote by $A_1, A_2, \dots, A_n$ the crossings of $\tilde{a}$ with $\tilde{b}$. We order crossings according to orientation of $\tilde{a}$. Similarly, we denote by $B_1, B_2, \dots, B_n$ the crossings of $\tilde{b}$ with $\tilde{a}$ when travelling from B_0 to B_∞ . The permutation $(\sigma(1), \sigma(2), \dots, \sigma(n))$ of $(1, 2, \dots, n)$ is defined by $A_i = B_{\sigma(i)}$ for each i . By A_i^- (resp. A_i^+) we denote a point on $\hat{a}$ or $\tilde{a}$ just before (resp. after) the crossing A_i with respect to the orientation of $\hat{a}$ or $\tilde{a}$. Note that both A_i^- and A_i^+ are not crossings of the projection or diagram. The points B_i^- and B_i^+ on $\hat{b}$ or $\tilde{b}$ are defined analogously. For points P and Q on a tangle projection $\hat{T}$ that are possibly but not necessarily crossings we denote by PQ the immersed arc in $\hat{T}$ starting from P and ending at Q with respect to the orientation of $\hat{T}$. In the case that both of them are mixed crossings there are two such possibilities. Then we specify the component. In case that both of them are self-crossings of the same component there may be two such possibilities. Then we take another point R and denote it by PRQ . See Fig. 2.7.




FIGURE 2.7.

The *vertical symmetry* of a 2-string tangle diagram is obtained by π -rotation along the vertical y -axis through the center of $\mathbb{B}^2$. Similarly a *horizontal symmetry* is obtained by π -rotation along the horizontal x -axis through the center of $\mathbb{B}^2$. We also use $\pi/2$ - (counter-clockwise) rotation. We denote the image of the vertical symmetry (resp. horizontal symmetry, $\pi/2$ -rotation) of a tangle T by $V(T)$ (resp. $H(T)$, $R(T)$). See Fig. 2.8.




FIGURE 2.8.

A *flying* (or *Tait flying*) is a local move of a link or tangle diagram (resp. projection) illustrated in Fig. 2.9 where $\tilde{S}$ (resp. $\hat{S}$) is any subdiagram (resp. subprojection).

If $\tilde{L}$ (resp. $\tilde{T}$) is obtained from $\tilde{L}_0$ (resp. $\tilde{T}_0$) by a series of flyings, then we say that $\tilde{L}$ (resp. $\tilde{T}$) is a *flying* of $\tilde{L}_0$ (resp. $\tilde{T}_0$).

Throughout the paper we use notation describing various kinds of twists performed on a diagram as illustrated in Fig. 2.10 and Fig. 2.11 and on a projection as illustrated in Fig. 2.12 (we code also in the notation the number or character of twists). Note that the direction of twists in this paper is opposite to the standard Conway's convention. We also remark that n is a variable so that even in the same




FIGURE 2.9.

figure n may be different. See for example Fig. 2.13, where a box can denote any positive even number.




FIGURE 2.10.




FIGURE 2.11.

By $T_1 + T_2$ we denote the sum of tangles T_1 and T_2 as illustrated in Figure 2.14. Sum of tangle diagrams and sum of tangle projections are similarly defined. We also use notations $T(n)$, $T(1/n)$, $\tilde{T}(n)$, $\tilde{T}(1/n)$, $\hat{T}(n)$ and $\hat{T}(1/n)$ where n is an integer as illustrated in Figure 2.14. Note that the notation $T(1/n)$ and $\tilde{T}(1/n)$ is used only for $n \geq 2$ to avoid the confusion that $T(1/1) \neq T(1)$ and $\tilde{T}(1/1) \neq \tilde{T}(1)$. By $T(a, b)$ we denote the sum $T(a) + T(b)$. This notation is based on Conway's tangle notation, but differs from it slightly. See Figure 2.14.

For a crossing P of a link diagram $\tilde{L}$, we say that the tangle diagram obtained by deleting a small disk centered at P from $\mathbb{S}^2$ is the *complementary tangle diagram* of $\tilde{L}$ at P .




FIGURE 2.12.




FIGURE 2.13. Every box can be replaced by an even positive number of twists

For a 2-string tangle T we define the *denominator* $D(T)$, *numerator* $N(T)$, X_+ -closure $X_+(T)$ and X_- -closure $X_-(T)$ as illustrated in Fig. 2.15.

Let $\tilde{L}$ (resp. $\tilde{T}$) be a positive diagram representing a link L (resp. tangle T). If the underlying projection $\hat{L}$ (resp. $\hat{T}$) is an underlying projection of another link L_0 (resp. tangle T_0), then clearly $L \geq L_0$ (resp. $T \geq T_0$). We use this fact throughout the paper without mentioning it explicitly. For the convenience we sometimes do not distinguish a link (resp. tangle) from its diagram and we will write for example that $\tilde{L} \geq L_0$ meaning that the link represented by $\tilde{L}$ is greater than or equal to L_0 .




FIGURE 2.14.




FIGURE 2.15.

2.2. Fundamental technique. The fundamental technique used in this paper is to apply descending (or ascending) algorithm that simplify or trivialize the link or tangle diagrams. As is well known in knot theory any link projection becomes a trivial link diagram if we give over/under crossing information according to descending algorithm. Namely we choose an order of components and choose a base point at each component and trace the link projection starting from the base point in chosen order. We then give over/under crossing information so that we meet each crossing as an over-crossing first time. We use descending algorithm to trivialize a part of the diagram often without explicitly mentioning it.

Many of the lemmas in this section follow closely that of [T-1]. We review them for completeness as their proofs illustrate our main method.

Lemma 2.1. *Let $\hat{L}$ (resp. $\hat{T}$) be a link (resp. tangle) projection. Let P be a self-crossing of a component $\hat{\ell}$ (resp. $\hat{t}$). Then $\hat{L} \geq r(\hat{L}, P)$ (resp. $\hat{T} \geq r(\hat{T}, P)$).*

Proof. Let $\tilde{L}$ be a diagram of a link L whose underlying projection is $r(\hat{L}, P)$. Let $\tilde{L}'$ be a diagram whose underlying projection is $\hat{L}$ with the following crossing information. For the crossings of $r(\tilde{L}, P)$ it is same as $\tilde{L}$, and for the crossings between $s(\tilde{L}, P)$ and $r(\tilde{L}, P)$, $s(\tilde{L}, P)$ is always over $r(\tilde{L}, P)$, and for the crossings of $s(\tilde{L}, P)$, it is determined by the descending algorithm so that $s(\tilde{L}, P)$ itself is unknotted. Then it is clear that $\tilde{L}'$ is also a diagram of L as desired. The tangle case is entirely the same and we omit it. $\square$

The following lemma is a diagram version of Lemma 2.1.

Lemma 2.2. *Let $\tilde{L}$ (resp. $\tilde{T}$) be a link (resp. tangle) diagram. Let P be a self-crossing of a component $\tilde{\ell}$ (resp. $\tilde{t}$). Suppose that the following conditions (1) and (2) hold.*

(1) *The knot represented by $s(\tilde{\ell}, P)$ (resp. $s(\tilde{t}, P)$) is greater than or equal to the trivial knot.*

(2) *Either $s(\tilde{\ell}, P)$ (resp. $s(\tilde{t}, P)$) is over $r(\tilde{\ell}, P)$ (resp. $r(\tilde{t}, P)$) at every negative crossings between them, or $s(\tilde{\ell}, P)$ (resp. $s(\tilde{t}, P)$) is under $r(\tilde{\ell}, P)$ (resp. $r(\tilde{t}, P)$) at every negative crossings between them.*

Then $\tilde{L} \geq r(\tilde{L}, P)$ (resp. $\tilde{T} \geq r(\tilde{T}, P)$).

Proof. Suppose that $s(\tilde{\ell}, P)$ is over $r(\tilde{\ell}, P)$ at every negative crossings between them. Then by changing some positive crossings between them to negative crossings we have a diagram $\tilde{L}'$ in which $s(\tilde{\ell}, P)$ is over $r(\tilde{\ell}, P)$ at every crossings. Note that $\tilde{L} \geq \tilde{L}'$. Then we have that $\tilde{L}'$ is a diagram of a link that is a connected sum of a link represented by $r(\tilde{L}, P)$ and a knot represented by $s(\tilde{\ell}, P)$. Then by the condition (1) we have that $\tilde{L}' \geq r(\tilde{L}, P)$. The other case is entirely analogous. $\square$

Lemma 2.3. *Let $\hat{L}_1$ and $\hat{L}_2$ (resp. $\hat{T}_1$ and $\hat{T}_2$) be a link (resp. tangle) projections that differ locally as illustrated in Fig 2.16 where P is a nugatory crossing of $\hat{L}_1$ (resp. $\hat{T}_1$) and R is a tangle (resp. subangle) possibly with some closed components. Then $\text{LINK}(\hat{L}_1) = \text{LINK}(\hat{L}_2)$ (resp. $\text{TANGLE}(\hat{T}_1) = \text{TANGLE}(\hat{T}_2)$).*

Proof. Let $\tilde{L}_1$ (resp. $\tilde{T}_1$) be a diagram whose underlying projection is $\hat{L}_1$ (resp. $\hat{T}_1$). Then by rotating the tangle (resp. subangle) $\tilde{R}$ we eliminate the nugatory crossing P and obtain the diagram $\tilde{L}_2$ (resp. $\tilde{T}_2$) of the same link (resp. tangle) whose underlying projection is $\hat{L}_2$ (resp. $\hat{T}_2$). Thus we have $\text{LINK}(\hat{L}_1) \subset \text{LINK}(\hat{L}_2)$ (resp. $\text{TANGLE}(\hat{T}_1) \subset \text{TANGLE}(\hat{T}_2)$). The converse holds in the same way. We can summarize the proof succinctly: P is a nugatory crossing for any $\tilde{L}_1$ (resp. $\tilde{T}_1$) with underlying projection $\hat{L}_1$ (resp. $\hat{T}_1$). $\square$

Lemma 2.4. *Let $\hat{T}_1$ and $\hat{T}_2$ be tangle projections that differs locally as illustrated in Fig. 2.17. Then $\hat{T}_1$ is greater than or equal to $\hat{T}_2$.*

Proof. Let $\tilde{T}_2$ be a diagram whose underlying projection is $\hat{T}_2$. Let $\tilde{T}_1$ be a diagram whose underlying projection is $\hat{T}_1$ such that the left string is over the right string in Fig. 2.17 and other crossings have the same over/under crossing information as $\tilde{T}_2$. Then $\tilde{T}_1$ and $\tilde{T}_2$ are transformed into each other by a second Reidemeister move. Therefore the result is proved. $\square$

Similarly we have the following lemma.




FIGURE 2.16.




FIGURE 2.17.

Lemma 2.5. *Let $\hat{T}_1$ and $\hat{T}_2$ be tangle projections that differs locally as illustrated in Fig. 2.18. Then $\hat{T}_1$ is greater than or equal to $\hat{T}_2$.*




FIGURE 2.18.

The proof is quite similar and we omit it.

In [P-1] and [T-1] it is shown that a knot projection $\hat{K}$ is a projection of a trefoil knot if and only if $\hat{K}$ is not almost trivial, and the link projection $\hat{L}$ is a projection of a Hopf link (plus, possibly, trivial components) if and only if $\hat{L}$ has mixed crossings (compare also [Co-G]). For tangle projections the analogous results hold:

Lemma 2.6. *A 1-string tangle projection $\hat{T}$ is a projection of the tangle in Fig. 2.19 (we call it a local trefoil) if and only if $\hat{T}$ is not an almost trivial tangle.*




FIGURE 2.19.

Proof. By Lemma 2.3 we may assume that $\hat{T}$ has no nugatory crossings. We trace $\hat{T}$ starting from its end point and then find a tear drop disk (1-gon) δ with vertex P , as illustrated in Fig. 2.20 (a). Then we continue to trace $\hat{T}$ and since P is not a nugatory crossing we return to δ at a crossing, say Q . See Fig. 2.20 (b). Then we add over/under crossing information to $\hat{T}$ by descending algorithm (except Q), as illustrated in Fig. 2.20 (c). $\square$




FIGURE 2.20.

Lemma 2.7. *A 2-string tangle projection $\hat{T}$ with vertical connection is a projection of the tangle $T(-2)$ (Fig. 2.21) (and its mirror image $T(2)$) if and only if $\hat{T}$ has mixed crossings.*




FIGURE 2.21.

Proof. Let A_1 be the first crossing on $\hat{a}$ between arcs $\hat{a}$ and $\hat{b}$. In particular the interior of A_0A_1 is disjoint from $\hat{b}$. We give over/under crossing information to A_1 so that it becomes a positive crossing. There are two cases depending on which string is over the other at the positive crossing A_1 . We use the notation A_i^- for a point on the arc $\hat{a}$ just before the crossing A_i and A_i^+ for a point just after the crossing. Neither A_i^- nor A_i^+ are crossings of $\hat{T}$. We give over/under crossing information to all other crossings of the tangle projection as follows.

- (i) If $\hat{a}$ is made to be under $\hat{b}$ at A_1 then we add over/under crossing information to all other crossings between arcs $\hat{a}$ and $\hat{b}$ in such a way that $\tilde{a}$ is over $\tilde{b}$ furthermore $\tilde{a}$ and $\tilde{b}$ are descending. The key point here is that a and b can be now deformed in two steps: first we deform a so it looks like on Fig. 2.22(b), in particular, a has only two crossings with b , one is A_1 and the other is near A_1 . Then we deform b to obtain the tangle $T(-2)$. See Fig. 2.22.
- (ii) If $\hat{a}$ is over $\hat{b}$ at A_1 then we add over/under crossing information to all other crossings so that $A_1^+A_\infty$ is under $\hat{b}$ and both $\tilde{a}$ and $\tilde{b}$ are descending. The key point here is that we first deform the string a Fig. 2.22(b), so that a has only two crossings with b , one is A_1 and the other is near A_1 . Then we deform b to obtain the tangle $T(-2)$. See Fig. 2.23.

□




FIGURE 2.22.




FIGURE 2.23.

Lemma 2.8. Let $\hat{T} = \hat{a} \cup \hat{b}$ be a 2-string tangle projection with vertical connection. Suppose that $\hat{T}$ has mixed crossings and the spine $\hat{a}'$ does not intersect $\hat{b}$. Then $\hat{T}$ is a projection of the tangle $T(1, 2)$ (Fig. 2.24).




FIGURE 2.24.

Proof. Let A_1 be the first mixed crossing of $\hat{T}$ on $\hat{a}$. By virtue of Lemma 2.1 we may assume that A_0A_1 has no self-crossings. Since A_1 is not on the spine $\hat{a}'$ we find, again by using Lemma 2.1, a tear drop disk δ as illustrated in Fig. 2.25 (a) or (b). Let B_i be the first mixed crossing in $\partial\delta$ on $\hat{b}$. Note that B_i may or may not equal to A_1 . Again by Lemma 2.1 we may assume that B_0B_i is simple. Thus we have the situation illustrated in Fig. 2.25 (a) or (b). In case (a) it is easy to see that $\hat{T}$ is a projection of $T(1, 2)$. In case (b) we add over/under crossing information so that the dotted line part of $\hat{a}$ is under everything and the dotted line part of $\hat{b}$ is under the real line parts of $\hat{a}$ and $\hat{b}$. Let P be the vertex of the 1-gon δ . Since $\hat{b}$ does not intersect A_0P we can deform $\hat{b}$ after deforming $\hat{a}$ into a neighborhood of $\partial B \cup A_0P$ as illustrated in Fig. 2.25 (b). $\square$




FIGURE 2.25.

Lemma 2.9. *A prime 2-string tangle projection $\hat{T}$ with a vertical connection is a projection of the tangle $T(1, 2)$ (Fig. 2.24) if and only if $\hat{T}$ is not the projection $\hat{T}(n)$ (see Fig 2.14) for any even n .*

Proof. Only tangles $T(m)$ with $|m| \leq n$ has $\hat{T}(n)$ as a projection. It is well known that the tangle $T(1, 2)$ is not $T(m)$ for any m . Thus we have proved the “only if” part. We will show the “if” part. First we note that the tangle $T(1, 2)$ has the vertical symmetry $T(1, 2) = V(T(1, 2)) = T(2, 1)$. Let $\hat{T} = \hat{a} \cup \hat{b}$ be a prime 2-string tangle projection with vertical connection that is not $\hat{T}(n)$ for any n . Let $\hat{b}'$ be the spine of $\hat{b}$. By Lemma 2.1 we have that $\hat{T}' = \hat{a} \cup \hat{b}'$ is a minor of $\hat{T}$. Suppose that $\hat{T}'$ has mixed crossings. Let $A_1, A_2, \dots, A_{2n}$ be the mixed crossings of $\hat{T}'$ that appear in this order on $\hat{a}$. Let $B_1, B_2, \dots, B_{2n}$ be the order of them on $\hat{b}'$ and σ the permutation $(\sigma(1), \sigma(2), \dots, \sigma(2n))$ of $(1, 2, \dots, 2n)$ defined by $A_i = B_{\sigma(i)}$ for each

i as before. We will show that $\hat{T}'$ is a projection of $T(1, 2)$ unless σ is an identical permutation. First suppose that $\sigma(2i-1) > \sigma(2i)$ for some $i \in \{1, 2, \dots, n\}$. Then we add over/under crossing information to $\hat{T}'$ as follows.

- (i) $A_0 A_{2i-1}^-$ is over everything.
- (ii) $A_{2i}^+ A_\infty$ is under everything.
- (iii) $\hat{a}$ is descending.
- (iv) $\hat{a}$ is under $\hat{b}'$ at A_{2i-1} and over $\hat{b}'$ at A_{2i} .

Then we have the tangle $T(1, 2)$. See Fig. 2.26 (a).

Next suppose that $\sigma(2j-1) < \sigma(2j)$ for each $j \in \{1, 2, \dots, \}$ and $\sigma(2i) > \sigma(2i+1)$ for some $i \in \{1, 2, \dots, n-1\}$. We take smallest such i . We further divide this case into the following two cases.

Case 1. $\sigma(2i+1) < \sigma(1)$. In this case we give over/under crossing information to $\hat{T}'$ as follows.

- (i) $A_1^+ A_{2i+1}^-$ is over everything.
- (ii) $A_{2i+1}^- A_\infty$ is under everything.
- (iii) Each of $A_0 A_1$, $A_1^+ A_{2i+1}^-$ and $A_{2i+1}^- A_\infty$ are descending.
- (iv) $\hat{a}$ is under $\hat{b}'$ at A_1 .

Then we have the tangle $T(1, 2)$. See Fig. 2.26 (b).

Case 2. $\sigma(2i+1) > \sigma(1)$. In this case we give over/under crossing information to $\hat{T}'$ as follows.

- (i) $A_1^+ A_{2i}^-$ is under everything.
- (ii) $A_{2i+1}^+ A_\infty$ is over everything.
- (iii) $A_0 A_1^-$ is under $A_{2i}^+ A_{2i+1}^-$.
- (iv) Each of $A_0 A_1$, $A_1^+ A_{2i}^-$, $A_{2i}^+ A_{2i+1}^-$ and $A_{2i+1}^+ A_\infty$ are descending.
- (v) $\hat{a}$ is over $\hat{b}'$ at A_1 and A_{2i} , and under $\hat{b}'$ at A_{2i+1} .

Then we have the tangle $T(1, 2)$. See Fig. 2.26 (c).




FIGURE 2.26.

Thus the rest is the case that $\sigma(i) < \sigma(i+1)$ holds for every $i \in \{1, 2, \dots, 2n-1\}$. This implies that $\sigma(i) = i$ for every i . Suppose that $A_i A_{i+1}$ intersects $A_{i+2j} A_{i+2j+1}$

for some i and $j > 0$ where we consider $A_{2n+1} = A_\infty$. Then by the use of Lemma 2.7 we have that $\hat{T}'$ is a projection of the tangle $T(1, 2)$. See for example Fig. 2.27.




FIGURE 2.27.

Thus we have the situation that there are mutually disjoint and possibly trivial 1-string subtangle projections $\hat{S}_1, \hat{S}_2, \dots, \hat{S}_{2n+1}$ such that the tangle $\hat{T}'$ is as illustrated in Fig. 2.28 (a). Note that in this situation it holds that for any self crossing P of $\hat{a}$ $s(\hat{a}, P)$ does not intersect $\hat{b}'$. Now we consider the spine $\hat{a}'$ and consider the tangle $\hat{T}'' = \hat{a}' \cup \hat{b}$. Since the tangle $T(1, 2)$ has the vertical symmetry as we have remarked at the beginning of the proof the same argument works for $\hat{T}''$ and we have the situation that there are mutually disjoint and possibly trivial 1-string subtangle projections $\hat{U}_1, \hat{U}_2, \dots, \hat{U}_{2n+1}$ such that the tangle $\hat{T}''$ is as illustrated in Fig. 2.28 (b). Since for any self-crossing P of $\hat{b}$ $s(\hat{b}, P)$ does not intersect $\hat{a}'$ we have that the set of mixed crossings of $\hat{T}''$ is exactly the same as the set of mixed crossings of $\hat{T}'$. Since $\hat{T}$ is prime and not equal to $\hat{T}(n)$ we have that at least one of $\hat{S}_1, \hat{S}_2, \dots, \hat{S}_{2n+1}$ has a crossing and then it must intersect at least one of $\hat{U}_1, \hat{U}_2, \dots, \hat{U}_{2n+1}$. Then by the use of Lemma 2.7 or Lemma 2.8 we obtain the tangle $T(1, 2)$ as illustrated in Fig. 2.29. $\square$

Lemma 2.10. *A prime 2-string tangle projection $\hat{T}$ with a X-connection is a projection of the tangle $T(1/3)$ (Fig. 2.30) if and only if $\hat{T}$ is not the projection $\hat{T}(n)$ (see Fig 2.14) for any odd n .*

Proof. The proof is quite similar to that of Lemma 2.9. Since the tangle $T(1/3)$ is different from any of the tangle $T(m)$, the “only if” part is proved. We will show the “if” part. Let $\hat{T} = \hat{a} \cup \hat{b}$ be a prime 2-string tangle projection with X-connection that is not $\hat{T}(n)$ for any n . Let $\hat{b}'$ be the spine of $\hat{b}$. Then $\hat{T}' = \hat{a} \cup \hat{b}'$ is a minor of $\hat{T}$. As in the proof of Lemma 2.9 we have that $\hat{T}'$ is a projection of $T(1/3)$ unless the permutation σ on $\{1, 2, \dots, 2n+1\}$ is the identity. See Fig. 2.31.

Thus the rest of the proof is the analysis of the case that $\sigma(i) = i$ for every i . Suppose that $A_i A_{i+1}$ intersects $A_{i+2j} A_{i+2j+1}$ for some i and $j > 0$ where we




FIGURE 2.28.




FIGURE 2.29.




FIGURE 2.30.




FIGURE 2.31.

consider $A_{2n+1} = A_\infty$. Then by the use of Lemma 2.7 we have that $\hat{T}'$ is a projection of the tangle $T(1/3)$. See for example Fig. 2.32.

Thus we have the situation that $\hat{T}'$ and $\hat{T}'' = \hat{a}' \cup \hat{b}$ is as illustrated in Fig. 2.33.




FIGURE 2.32.




FIGURE 2.33.

Then by the use of Lemma 2.7 or Lemma 2.8, we have the tangle $T(1/3)$ as illustrated in Fig. 2.34. $\square$




FIGURE 2.34.

Lemma 2.11. *Let $\hat{T}$ be a prime 2-string tangle projection with vertical connection. Suppose that the right string $\tilde{b}$ has no self-crossings and $\hat{T}$ is not a projection of the tangle $T(-4)$ (Fig. 2.35). Then one of the following (1) or (2) holds.*

- (1) *There are a tangle projection $\hat{S}$ with horizontal connection, and natural numbers m and n such that $\hat{T} = \hat{S} + R(\hat{T}(2m-1, 2n-1))$ (Fig. 2.36 (a)).*
- (2) *There is a tangle projection $\hat{S}$ with X-connection and a natural number m such that $\hat{T} = \hat{S} + \hat{T}(2m)$ (Fig. 2.36 (b)).*

Proof. We analyze a permutation $(\sigma(1), \sigma(2), \dots, \sigma(2k))$ of $(1, 2, \dots, 2k)$ defined by $A_i = B_{\sigma(i)}$. Note that the tangle projection in Fig. 2.36 (b) defines the “backward”




FIGURE 2.35.




FIGURE 2.36.

permutation $\sigma(2m) < \sigma(2m-1) < \dots < \sigma(2) < \sigma(1)$. Namely $\sigma(i) = 2m+1-i$ for every i . The tangle projection in Fig. 2.36 (a) defines the permutation composed of two “backward” pieces, that is, $\sigma(2n-1) < \sigma(2n-2) < \dots < \sigma(1) < \sigma(2m+2n-2) < \sigma(2m+2n-3) < \dots < \sigma(2n+1) < \sigma(2n)$. We will show that $\hat{T}$ is a projection of $T(-4)$ unless the permutation σ is one of these forms. For that purpose we watch the “forward” part of σ . First suppose that $\sigma(i) > \sigma(i+1)$ for every i . Then we have $\sigma(i) = 2k+1-i$ for each i . Next suppose that $\sigma(2i) < \sigma(2i+1)$ for some i . In this case we give over/under crossing information to $\hat{T}$ as follows.

- (i) $A_0 A_{2i}^-$ is under everything.
- (ii) $A_{2i+1}^+ A_\infty$ is over everything.
- (iii) $\hat{a}$ is descending.
- (iv) $\hat{a}$ is over $\hat{b}$ at A_{2i} and under $\hat{b}$ at A_{2i+1} .

Then we have $T(-4)$ as illustrated in Fig. 2.37

Next suppose that $\sigma(2j) > \sigma(2j+1)$ for every j and there is i such that $\sigma(2i-1) < \sigma(2i)$. We fix this i and analyze how it goes after A_{2i} (Case 1, Case 2) and how it goes back before A_{2i-1} (Case 1', Case 2') .

Case 1. There exists $j > i$ such that $\sigma(2i-1) < \sigma(2j-1) < \sigma(2j)$.

We take the smallest such j . Then by the assumption we have $\sigma(2j-1) < \sigma(2j-2) < \dots < \sigma(2i)$. In this case we give over/under crossing information to $\hat{T}$ as follows.

- (i) $A_0 A_{2i-1}^-$ is over everything.
- (ii) $A_{2j}^+ A_\infty$ is under everything.
- (iii) $\hat{a}$ is descending.




FIGURE 2.37.

- (iv) $\hat{a}$ is under $\hat{b}$ at A_{2i-1} and A_{2j-1} and over $\hat{b}$ at $A_{2i}, A_{2i+1}, \dots, A_{2j-2}$ and A_{2j} .

Then we have $T(-4)$ as illustrated in Fig. 2.38




FIGURE 2.38.

Case 1'. There exists $l < i$ such that $\sigma(2i) > \sigma(2l) > \sigma(2l-1)$.

We take the largest such l . Then by the assumption we have $\sigma(2l) > \sigma(2l+1) > \dots > \sigma(2i-1)$. Note that this case is obtained from Case 1 by reversing the order. Namely $2i-1, 2i, 2j-1$ and $2j$ correspond to $2i, 2i-1, 2l$ and $2l-1$ respectively. Since the tangle $T(-4)$ has horizontal symmetry $H(T(-4)) = T(-4)$ we have that $\hat{T}$ is a projection of $T(-4)$.

Case 2. Case 1 does not occur and there exist $j > i$ such that $\sigma(2j) < \sigma(2i-1) < \sigma(2j-1)$.

We take the smallest such j . Then by the assumption we have $\sigma(2j-1) < \sigma(2j-2) < \dots < \sigma(2i)$. In this case we give over/under crossing information to $\hat{T}$ as follows.

- (i) $A_0 A_{2i-1}^-$ is under everything.
- (ii) $A_{2j}^+ A_\infty$ is over everything.
- (iii) $A_0 A_{2j}$ and $A_{2j} A_\infty$ are descending.
- (iv) $\hat{a}$ is under $\hat{b}$ at A_{2i-1} and A_{2j-1} and over $\hat{b}$ at $A_{2i}, A_{2i+1}, \dots, A_{2j-2}$ and A_{2j} .

Then we have $T(-4)$ as illustrated in Fig. 2.39




FIGURE 2.39.

Case 2'. Case 1' does not occur and there exist $l < i$ such that $\sigma(2l-1) > \sigma(2i) > \sigma(2l)$.

By the horizontal symmetry of $T(-4)$ and by Case 2 we have that $\hat{T}$ is a projection of $T(-4)$.

In the following cases we assume that neither Case 1 nor Case 1', Case 2 or Case 2' occurs.

Then after A_{2i} we have the following two possibilities.

Case P1. There exists $j > i$ such that $\sigma(2j-1) < \sigma(2i-1) < \sigma(2j-2)$.

In this case we take the smallest such j and therefore we further have $\sigma(2j-1) < \sigma(2j-2) < \dots < \sigma(2i)$ (or $\sigma(2j-1) < \sigma(2j-2) = \sigma(2i)$ as a special case).

Case P2. Case P1 does not occur. Namely $\sigma(2i-1) < \sigma(2k) < \sigma(2k-1) < \dots < \sigma(2i)$ (or $2k = 2i$ as a special case).

Before A_{2i-1} we have the following two possibilities.

Case Q1. There exists $l < i$ such that $\sigma(2l) > \sigma(2i) > \sigma(2l-2)$.

In this case we take the largest such l and therefore we further have $\sigma(2l) > \sigma(2l+1) > \dots > \sigma(2i-1)$ (or $\sigma(2l) > \sigma(2l+1) = \sigma(2i-1)$ as a special case).

Case Q2. Case Q1 does not occur. Namely $\sigma(2i) > \sigma(1) > \sigma(2) > \dots > \sigma(2i-1)$ (or $1 = 2i-1$ as a special case).

Thus we have the following four cases to be considered.

Case P1Q1. In this case we give over/under crossing information to $\hat{T}$ as follows.

- (i) $A_0 A_{2l}^-$ is under everything.
- (ii) $A_{2j-1}^+ A_\infty$ is over everything.
- (iii) $A_{2l}^+ A_{2i-1}^-$ is over everything except $A_{2j-1}^+ A_\infty$.
- (iv) $A_{2i}^+ A_{2j-1}^-$ is under everything except $A_0 A_{2l}^-$.
- (v) Each of $A_0 A_{2l}$, $A_{2l} A_{2i-1}$, $A_{2i-1} A_{2i}$, $A_{2i} A_{2j-1}$ and $A_{2j-1} A_\infty$ is descending.
- (vi) $\hat{a}$ is under $\hat{b}$ at A_{2l} , A_{2i-1} , $A_{2i+1} A_{2i+2}$, $\dots$, A_{2j-1} and over $\hat{b}$ at A_{2l+1} , A_{2l+2} , $\dots$, A_{2i-2} and A_{2i} .

Then we have $T(-4)$ as illustrated in Fig. 2.40

Case P1Q2. In this case we give over/under crossing information to $\hat{T}$ as follows.

- (i) $A_{2j-1}^+ A_\infty$ is over everything.
- (ii) Each of $A_0 A_{2j-1}$ and $A_{2j-1} A_\infty$ is descending.
- (iii) $\hat{a}$ is over $\hat{b}$ at A_1 , A_2 , $\dots$, A_{2i-2} and A_{2i} and under $\hat{b}$ at A_{2i+1} , A_{2i+2} , $\dots$, A_{2j-1} .

Then we have $T(-4)$ as illustrated in Fig. 2.41




FIGURE 2.40.




FIGURE 2.41.

Case P2Q1. This case follows from Case P1Q2 and the horizontal symmetry of $T(-4)$.

Case P2Q2. Suppose that $\sigma(1) > \sigma(2k)$. Then we give over/under crossing information to $\hat{T}$ as follows.

- (i) $A_1^+ A_{2i}^-$ is over everything.
- (ii) $A_{2i}^+ A_{2k}^-$ is under everything.
- (iii) Each of $A_0 A_1$, $A_1 A_{2i}$, $A_{2i} A_{2k}$ and $A_{2k} A_\infty$ is descending.
- (iv) $A_0 A_1$ is under $A_{2k} A_\infty$.
- (v) $\hat{a}$ is over $\hat{b}$ at A_{2i} and A_{2k} and under $\hat{b}$ at A_1 .

Then we have $T(-4)$ as illustrated in Fig. 2.42

Thus we finally have that $\sigma(2i-1) < \sigma(2i-2) < \dots < \sigma(1) < \sigma(2k) < \sigma(2k-1) < \dots < \sigma(2i+1) < \sigma(2i)$. This is one of the permutations described at the beginning of the proof.

Now suppose that $\sigma(2i-1) > \sigma(2i)$ for every i . Namely there are no forward part in σ and so we have $\sigma(2k) < \sigma(2k-1) < \dots < \sigma(2) < \sigma(1)$. This is the second permutation described at the beginning of the proof. Suppose that in these cases there are i and j with $i < j$ such that $A_i^+ A_{i+1}^-$ intersects $A_j^+ A_{j+1}^-$. Then by the use of Lemma 2.7 we have $T(-4)$ as illustrated in Fig. 2.43. Then by the primeness of $\hat{T}$ we reach the desired conclusion. $\square$

The following corollaries of Lemma 2.11 are the main tools of this section.




FIGURE 2.42.




FIGURE 2.43.

Corollary 2.12. *Let $\hat{T}$ be a prime 2-string tangle projection with vertical connection. Suppose that the right string $\hat{b}$ has no self-crossings and $\hat{T}$ is not a projection of the tangle $T(-4)$, $R(T(-1, -2)) + T(-2)$, or $T(-3, 2)$ (Fig. 2.44). Then one of the followings holds.*

- (1) $\hat{T} = \hat{T}(1/2l) + R(\hat{T}(2m - 1, 2n - 1))$ (Fig. 2.45 (a)) for some $l \geq 0$ and $m, n > 0$.
- (2) $\hat{T} = \hat{T}(1/(2l - 1), 2m)$ (Fig. 2.45 (b)) for some $l, m > 0$.




FIGURE 2.44.




FIGURE 2.45.

Proof. Assume that $\hat{T}$ is not a projection of the tangle $T(-4)$ then (1) or (2) of Lemma 2.11 holds.

First suppose that the condition (1) of Lemma 2.11 holds. Note that the tangle projection $\hat{S} + R(\hat{T}(1, 1)) = \hat{S} + \hat{T}(2)$ is a minor of $\hat{S} + R(\hat{T}(2m-1, 2n-1))$. We apply Lemma 2.9 to $\hat{S}$ and have that $\hat{S} + \hat{T}(2)$ is a projection of $R(T(-1, -2)) + T(-2)$ unless $\hat{S} = \hat{T}(1/2l)$ for some l . Thus we have the conclusion (1).

Next suppose that the condition (2) of Lemma 2.11 holds. Note that the tangle projection $\hat{S} + \hat{T}(2)$ is a minor of $\hat{S} + \hat{T}(2m)$. We apply Lemma 2.10 to $\hat{S}$ and have that $\hat{S} + \hat{T}(2)$ is a projection of $T(-3, 2)$ unless $\hat{S} = \hat{T}(1/(2l-1))$ for some l . Thus we have the conclusion (2). $\square$

Lemma 2.13. *Let $\hat{K}$ be a knot projection. Suppose that $\hat{K}$ is not an underlying projection of the $(2, 5)$ -torus knot. Then $\hat{K}$ is a connected sum of some knot projections each of which is one of the projections in Fig. 2.46.*

Proof. Suppose that $\hat{K}$ is a connected sum of prime projections $\hat{K}_1, \dots, \hat{K}_k$. Then each $\hat{K}_i$ is not an underlying projection of the $(2, 5)$ -torus knot. We will show that each $\hat{K}_i$ is one of the projections in Fig. 2.46. Suppose that $\hat{K}_i$ has two or more crossings. Since $\hat{K}_i$ is prime we have that $\hat{K}_i$ is reduced. Choose a 1-gon disk (tear drop disk) δ in $\hat{K}_i$ as follows: orient $\hat{K}_i$ and choose a basepoint P on $\hat{K}_i$ that is not a crossing point. Follow $\hat{K}_i$, starting from P , and let P_0 be the first point (crossing) at which one meets his trace. The simple closed curve which was traced divides the




FIGURE 2.46.

sphere of the projection into two disks and δ is the disk which does not contain P . Let $\hat{T}$ be the complementary tangle projection of the vertex P_0 . If $\hat{T}$ is a projection of the tangle $T(-4)$, $R(T(-1, -2)) + T(-2)$, or $T(-3, 2)$ then $\hat{K}_i$ is a projection of the $(2, 5)$ -torus knot as illustrated in Fig. 2.47. Then we have by Lemma 2.12 that $\hat{T} = \hat{T}(1/2l) + R(\hat{T}(2m - 1, 2n - 1))$ or $\hat{T} = \hat{T}(1/(2l - 1), 2m)$. Thus we have that $\hat{K}_i$ is one of the projections in Fig. 2.46. $\square$




FIGURE 2.47.

2.3. The main results on Positive Links.

Theorem 2.14. *Let $\hat{L} = \hat{K}_1 \cup \hat{K}_2 \cup \dots \cup \hat{K}_\mu$ be an oriented link projection. Then either*

- (A) *$\hat{L}$ is an underlying projection of the $(2, 5)$ -torus knot (Fig. 2.47) (with possible trivial components) or*
- (B) *Each $\hat{K}_i$ is a knot projection described in Lemma 2.13 and*
 - (1) *$\hat{L}$ is an underlying projection of the $(2, 4)$ -torus link (Fig. 1.4 (c)) (with possible trivial components) or*
 - (2) *$\hat{L}$ is an underlying projection of the $(3, 3)$ -torus link (Fig. 2.55 (a)) (with possible trivial components) or*
 - (3) *$\hat{L}$ is an underlying projection of 8_{10}^3 (Fig. 2.55 (b)) (with possible trivial components) or*
 - (4) *$\hat{L}$ is an underlying projection of a connected or disjoint sum of the right-handed trefoil knot and the right-handed Hopf link (with possible trivial components) or*
 - (5) *$\hat{L}$ is an underlying projection of a disjoint and/or connected sum of three right-handed Hopf links (with possible trivial components) or*
 - (6) *$\hat{L}$ has no mixed crossings or*

- (7) $\hat{L}$ is an $R1$ augmentation of $N(\hat{T}(p_1, p_2, p_3))$ with anti-parallel orientations of components for some positive even numbers p_1, p_2 and p_3 (Fig. 2.48) with possible trivial circles or
- (8) $\hat{L}$ is an $R1$ augmentation of $D(\hat{T}(2k))$ with anti-parallel orientation of components for some positive number k (Fig. 2.49) with possible trivial circles or a connected or disjoint sum of two copies of such link projections.




FIGURE 2.48.




FIGURE 2.49.

Proof. Suppose that some $\hat{K}_i$ is a projection of the $(2, 5)$ -torus knot. Then we are dealing with Case (A). Suppose that each $\hat{K}_i$ is not a projection of the $(2, 5)$ -torus knot. Then by Lemma 2.13 we have that $\hat{K}_i$ is a connected sum of some knot projections each of which is a projection in Fig. 2.46.

Suppose that $\hat{K}_i$ is not almost trivial and $\hat{K}_i$ has some mixed crossings. Let P be a mixed crossing on $\hat{K}_i$ with $\hat{K}_j$ where $\hat{K}_i$ crosses from left to right of $\hat{K}_j$. Then we add over/under crossing information to $\hat{K}_i \cup \hat{K}_j$ so that $\hat{K}_j$ produces a trivial knot, $\hat{K}_i$ produces a right-handed trefoil knot and $\hat{K}_j$ is over $\hat{K}_i$ except P . Then the result is a connected sum of a right-handed trefoil knot and a right-handed Hopf link. Thus we have case (B) (4). Suppose that $\hat{K}_i$ has no mixed crossings but some $\hat{K}_j$ and $\hat{K}_l$ has a common mixed crossing. Then by a similar argument we have that $\hat{L}$ is an underlying projection of a disjoint sum of a right-handed trefoil knot and a right-handed Hopf link. Namely we again have case (B) (5). Suppose that $\hat{L}$ has no mixed crossings. Then we have case (B) (6). Thus in the following we assume that each $\hat{K}_i$ is an almost trivial projection.

Now suppose that $\hat{K}_i$ and $\hat{K}_j$ has a common mixed crossing, say P . We see the complementary tangle projection $\hat{T}$ of $\hat{K}_i \cup \hat{K}_j$ at P . We apply Lemma 2.10 to $\hat{T}$

and have that $\hat{K}_i \cup \hat{K}_j$ is a projection of the $(2, 4)$ -torus link (Fig. 1.4 (c)) unless $\hat{K}_i \cup \hat{K}_j$ is an R1 augmentation of the link projection $D(\hat{T}(2n))$ with anti-parallel orientation for some natural number n (Fig. 2.49). If $\hat{K}_i \cup \hat{K}_j$ is a projection of the $(2, 4)$ -torus link then we have the case (B) (1). Thus we consider the case that for any $\hat{K}_i$ and $\hat{K}_j$ that has a common mixed crossing the union $\hat{K}_i \cup \hat{K}_j$ is an R1 augmentation of the link projection $D(\hat{T}(2n))$ with anti-parallel orientation for some natural number n .

Let $\hat{L} = \hat{L}_1 \cup \hat{L}_2 \cup \cdots \cup \hat{L}_u$ be the connected components of $\hat{L}$. Namely each $\hat{L}_i$ is a connected projection and $\hat{L}_i$ and $\hat{L}_j$ are disjoint for $i \neq j$.

Let $\mu(i)$ be the number of knot projections that compose $\hat{L}_i$. We may suppose without loss of generality that $\mu(1) \geq \mu(2) \geq \cdots \geq \mu(u)$. Suppose that $\mu(1) \geq \mu(2) \geq \mu(3) \geq 2$. Then for each $i = 1, 2, 3$, choose two knot projections, say $\hat{K}_a$ and $\hat{K}_b$ in $\hat{L}_i$ that has a mixed crossing, say P . We may suppose without loss of generality that $\hat{K}_b$ crosses from left to right of $\hat{K}_a$ at P . Then we add over/under crossing information to $\hat{L}_i$ such that $\hat{K}_a$ is over $\hat{K}_b$ except at P . We trivialize other parts and we have that each $\hat{L}_i$ is a projection of a right-handed Hopf link with possible trivial components. Thus we have case (B) (5). Next we consider the case that $\mu(3) = 1$ or $u \leq 2$. Suppose that $\mu(1) \geq 4$. Let $\hat{K}_{a_1}, \hat{K}_{a_2}, \dots, \hat{K}_{a_{\mu(1)}}$ be the knot projections in $\hat{L}_1$. Consider an abstract graph G with vertices $\hat{K}_{a_1}, \hat{K}_{a_2}, \dots, \hat{K}_{a_{\mu(1)}}$ and edges $\hat{K}_{a_i} \hat{K}_{a_j}$ where $\hat{K}_{a_i}$ and $\hat{K}_{a_j}$ have mixed crossings. Let H_0 be a spanning tree of G . By re-ordering the vertices if necessary we may suppose that K_{a_1} is a free vertex (degree one vertex) of H_0 . Let P_1 be a mixed crossing on $\hat{K}_{a_1}$ at which $\hat{K}_{a_1}$ crosses from right to left of the other component. Let H_1 be the graphs obtained from H_0 by removing the vertex K_{a_1} together with the edge incident to it. We may again suppose without loss of generality that K_{a_2} is a free vertex of H_1 . Let P_2 be a mixed crossing on $\hat{K}_{a_2}$ at which $\hat{K}_{a_2}$ crosses from right to left of the other component. Let H_2 be the graphs obtained from H_1 by removing the vertex K_{a_2} together with the edge incident to it. We continue this process and finally reach a graph $H_{\mu(1)-1}$ on a single vertex $K_{a_{\mu(1)}}$. Now we give over/under crossing information to $\hat{L}_1$ so that $\hat{K}_{a_i}$ is over $\hat{K}_{a_j}$ if $i < j$ except at $P_1, P_2, \dots, P_{\mu(1)-2}$ and $P_{\mu(1)-1}$. Then we have that the result is a connected sum of $\mu(1) - 1$ right-handed Hopf links whose linking number structure respects the tree H_0 . Thus the rest is the case that $2 \leq \mu(1) \leq 3$. Suppose that $\mu(1) = 3$ and $\mu(2) \geq 2$. Then by giving over/under crossing information to $\hat{L}$ in a similar way we have that $\hat{L}$ is a projection of the disjoint sum of two links, one is a connected sum of two right-handed Hopf links, and the other is a right-handed Hopf link. Namely we again have the case (B) (5).

Suppose that $\mu(1) = 3$ and $\mu(2) = 1$ or $u = 1$. We may suppose that $\hat{K}_1, \hat{K}_2$ and $\hat{K}_3$ are the components of $\hat{L}_1$. We may suppose without loss of generality that $\hat{K}_1$ and $\hat{K}_2$ have common mixed crossings and $\hat{K}_2$ and $\hat{K}_3$ have common mixed crossings. Since $\hat{K}_1 \cup \hat{K}_2$ is an R1 augmentation of the link projection $D(\hat{T}(2n))$ with anti-parallel orientation it is as illustrated in Fig. 2.50 where the tangles are almost trivial 1-tangles. Let $A_1, A_2, \dots, A_{2n}$ be the common mixed crossings of $\hat{K}_1$ and $\hat{K}_2$ that appears in this order on $\hat{K}_2$ along the orientation of $\hat{K}_2$.

Case 1. $\hat{K}_1$ and $\hat{K}_3$ have no common mixed crossings.




FIGURE 2.50.

First suppose that $\hat{K}_3$ has intersection with only one of $A_1A_2, A_2A_3, \dots, A_{2n-1}A_{2n}$. Then we have that the projection $\hat{L}_1$ is a connected sum of two link projections and we have the case (B) (8). Next suppose that $\hat{K}_3$ has intersection with at least two of $A_1A_2, A_2A_3, \dots, A_{2n-1}A_{2n}$, say A_1A_2 and $A_{2i-1}A_{2i}$. Since $\hat{K}_2 \cup \hat{K}_3$ is also an R1 augmentation of the link projection $D(\hat{T}(2m))$ with anti-parallel orientation for some natural number m it has the property that if we do smoothing at a self-crossing of $\hat{K}_2$ only one of the two components of the result has intersection with $\hat{K}_3$. This implies that $\hat{K}_3$ do not intersects the R1 residual of $\hat{K}_2$. Thus in this case we may assume that each of $\hat{K}_1, \hat{K}_2$ and $\hat{K}_3$ is a simple closed curve on $\mathbb{S}^2$. Then we have that $\hat{L}_1$ is as illustrated in Fig. 2.51 (a) and it is a projection of the link 8_{10}^3 . See for example Fig. 2.51 (b). Thus we have case (B) (3).

Case 2. $\hat{K}_1$ and $\hat{K}_3$ have common mixed crossings.

First suppose that $\hat{K}_3$ intersects R1 residual of $\hat{K}_1$ or $\hat{K}_2$, say $\hat{K}_2$. Then by taking a minor of $\hat{L}_1$ using Lemma 2.1 if necessary we have one of the situations illustrated in Fig. 2.52 and Fig. 2.53. Then we have that $\hat{L}_1$ is a projection of the $(3, 3)$ -torus link as is also illustrated in 2.52 and Fig. 2.53.

Thus we may assume that $\hat{K}_3$ has no intersection with R1 residuals of $\hat{K}_1$ and $\hat{K}_2$. Therefore we may assume that both $\hat{K}_1$ and $\hat{K}_2$ are simple closed curves on $\mathbb{S}^2$. By changing the role of $\hat{K}_2$ and $\hat{K}_3$ and doing the same argument we may further assume that $\hat{K}_1$ is also a simple closed curve on $\mathbb{S}^2$. Suppose that there are consecutive crossings on $\hat{K}_3$ each of which $\hat{K}_3$ crosses from left to right, or from right to left. Then we have that $\hat{L}_1$ is a projection of $(3, 3)$ -torus link as illustrated in Fig. 2.54 (a). Thus we have that two consecutive crossings P_1 and P_2 on $\hat{K}_3$ are placed as illustrated in Fig 2.54 (b). (Since $(3, 3)$ -torus link is invertible and $\hat{K}_1$ and $\hat{K}_2$ are exchangeable omit another orientation possibility.) If the next crossing




FIGURE 2.51.

P_3 on $\hat{K}_3$ is before P_2 with respect to the orientation of the other curve then we have the $(3, 3)$ -torus link as illustrated in Fig. 2.54 (c). Therefore we have the situation illustrated in Fig. 2.54 (d). Then we have that only possible projection is a projection in Fig. 2.48. See Fig. 2.54 (e). Thus we have case (B) (7).

Finally suppose that $\mu(1) = 2$. Then we have the case (B) (8). This completes the proof. $\square$

Theorem 2.15. *Let L be a nontrivial positive link. Then either*

- (A) $L \geq (2, 5)$ -torus knot (Fig. 2.47) (with possible trivial components) or
- (B) *Each component of L is a connected sum of pretzel knots $L(p_1, p_2, p_3)$ where p_1, p_2 and p_3 are positive odd numbers (Fig. 1.1) or the unknot and*
 - (1) $L \geq (2, 4)$ -torus link (Fig. 1.4 (c)) (with possible trivial components) or
 - (2) $L \geq (3, 3)$ -torus link (Fig. 2.55 (a)) (with possible trivial components) or
 - (3) $L \geq 8^3_{10}$ (Fig. 2.55 (b)) (with possible trivial components) or
 - (4) $L \geq$ connected or disjoint sum of the right-handed trefoil knot and the right-handed Hopf link (with possible trivial components) or
 - (5) $L \geq$ disjoint and/or connected sum of three right-handed Hopf links (with possible trivial components) or
 - (6) L is a (connected and/or disjoint sum of) pretzel knot(s) $L(p_1, p_2, p_3)$ (Fig. 1.1) (with possible trivial components) or
 - (7) L is a three-component pretzel link $L(p_1, p_2, p_3)$ (Fig. 2.55 (c)) (with possible trivial components) or
 - (8) L is a $(2, 2k)$ -torus link with anti-parallel orientation of components (Fig. 2.55 (d)) or the connected or disjoint sum of two copies of such links (with possible trivial components).

Proof. First note that a positive diagram whose underlying projection is the projection illustrated in Fig. 2.46 (d) represents a pretzel knot $L(1, p_2, p_3)$ for some positive odd numbers p_2 and p_3 . See Fig. 2.56. Also note that a pretzel knot $L(p_1, p_2, p_3)$ where p_1, p_2 and p_3 are positive odd numbers is greater than or equal to the right-handed trefoil knot $L(1, 1, 1)$. Let $\tilde{L}$ be a positive diagram of L and




FIGURE 2.52.

$\hat{L}$ its underlying projection. Then by Theorem 2.14 and the facts noted above we have the result. $\square$

Corollary 2.16. *Let L be a nontrivial positive link, then either $\sigma(L) \leq -3$ or:*

- (a) *L is a pretzel knot $L(p_1, p_2, p_3)$ for some positive odd numbers p_1, p_2 and p_3 (Fig. 1.1) (with possible trivial components); then $\sigma(L) = -2$ or*
- (b) *L is a three component pretzel link $L(p_1, p_2, p_3)$ for some positive even numbers p_1, p_2 and p_3 (Fig. 2.55 (c)) (with possible trivial components); then $\sigma(L) = -2$ or*
- (c) *L is a $(2, 2k)$ -torus link with anti-parallel orientation of components (Fig. 1.5) (with possible trivial components); then $\sigma(L) = -1$ or*




FIGURE 2.53.

- (d) L is a connected or disjoint sum of two copies of links from (c) (with possible trivial components); then $\sigma(L) = -2$.

In particular if K is a nontrivial positive knot, then either $\sigma(L) \leq -4$ or (a) holds.

Proof. It follows immediately from the fact that the signatures of the (2,5)-torus knot, (2,4)-torus link, (3,3)-torus link, and the link 8^3_{10} (Fig. 2.55 (b)) are -4,-3,-4 and -3 respectively, and that if $L_1 \geq L_2$ then, by Giller inequality, $\sigma(L_1) \leq \sigma(L_2)$. $\square$

Remark 2.17. We work, in the next section, with almost positive links and demonstrate there, in particular, the title result of the paper that a nontrivial Almost Positive Links have negative Signature (Corollary 1.7). However Corollary 1.7 can be also derived from Corollary 2.16. The outline of this derivation is as follows. Let $\tilde{L}$ be an almost positive diagram of a link L . Let $\tilde{L}_+$ be the positive diagram obtained from $\tilde{L}$ by changing the negative crossing to a positive crossing. Let L_+ be the link represented by $\tilde{L}_+$. If $\sigma(L_+) \leq -3$ then we have $\sigma(L) \leq -1 < 0$ as desired. Then by checking the exceptional cases (a)-(d) we obtain the result. In the next section




FIGURE 2.54.




FIGURE 2.55.

we give a different proof of Corollary 1.7. It applies Theorem 3.2 which is based on Lemma 3.1. This lemma plays an important role also in Section 4.

3. ALMOST POSITIVE LINKS

Lemma 3.1. *Let $\tilde{T}$ be a reduced almost positive 2-string tangle diagram with vertical connection. Suppose that the negative crossing is a mixed crossing. If $\tilde{T}$ is not*




FIGURE 2.56.

greater than or equal to any of the tangles $T(-2)$, $T(3_1, 0_1)$ and $T(0_1, 3_1)$ in Fig. 3.1, then $\tilde{T}$ is one of the diagrams $\tilde{T}_{1+}$, $\tilde{T}_{1-}$, $\tilde{T}_{2+}$ and $\tilde{T}_{2-}$ in Fig. 3.2.




FIGURE 3.1.




FIGURE 3.2.

Proof. Let N be the negative crossing of $\tilde{T}$. We have by Lemma 2.6 that both components of $\tilde{T}$ are almost trivial.

Case 1. $\tilde{T}$ has just two mixed crossings.

Then we have that these crossings are related on both strings and we have that $\tilde{T}$ is one of $\tilde{T}_{1\pm}$.

Case 2. $\tilde{T}$ has four or more mixed crossings.

Case 2.1. $\tilde{T}$ has a self-crossing. We may suppose without loss of generality that $\tilde{a}$ has a self-crossing.

First suppose that $m(N, \tilde{a}) \geq 1$ and P is the root of N on $\tilde{a}$. Suppose that the positive tangle $r(\tilde{T}, P)$ has mutual crossings. Then by Lemma 2.7 we have that $r(\tilde{T}, P) \geq T(-2)$. Since we have $\tilde{T} \geq r(\tilde{T}, P)$ by Lemma 2.2 we have $\tilde{T} \geq T(-2)$. Thus we may suppose that if $m(N, \tilde{a}) \geq 1$ and P is the root of N on $\tilde{a}$, then $r(\tilde{T}, P)$ has no mutual crossings. Then we have that N is rightmost. We have one of the following:

- Case 2.1.1. There are two positive mixed crossings that are related on $\tilde{a}$ which are rightmost and have multiplicity greater than zero on $\tilde{a}$.
Case 2.1.2. Case 2.1.1 does not hold and there are two positive crossings that are related on $\tilde{a}$ which are not rightmost.

In Case 2.1.1 we can choose such crossings B_i and B_j ($i < j$) so that i is odd and B_k is not related to B_i on $\tilde{a}$ for $i < k < j$.

Then we can operate, according to the position of N , one of the following two crossing changes without changing the negative crossing N .

- (1) $B_0 B_i^-$ is over the other parts and $B_j^+ B_\infty$ is under the other parts.
- (2) $B_0 B_i^-$ is under the other parts and $B_j^+ B_\infty$ is over the other parts.

Then we have $\tilde{T} \geq T(-2)$ as illustrated in Fig. 3.3.




FIGURE 3.3.

In Case 2.1.2, using also Lemma 2.7, we have $\tilde{T} \geq T(-2)$ as illustrated (up to horizontal symmetry) in Fig. 3.4.

Case 2.2. $\tilde{T}$ has no self-crossings.

Suppose that N is neither the first crossing nor the last crossing on $\tilde{b}$. Then we have one of the situations illustrated in Fig. 3.5 and then have $\tilde{T} \geq T(-2)$ as illustrated.




FIGURE 3.4.

Therefore N must be the first or the last crossing on $\tilde{b}$. In addition if there is a part in $\tilde{T}$ illustrated in Fig. 3.6 then we have $\tilde{T} \geq T(-2)$. Then an easy analysis forces $\tilde{T}$ to be $\tilde{T}_{\pm}$. This completes the proof. $\square$

In the following we call the technique used in Case 2.1.1 as “over and under technique”.

Theorem 3.2. *Let $\tilde{L}$ be a reduced almost positive diagram of a link L . If $\tilde{L}$ is not a diagram in Fig. 1.2 (plus trivial circles), then*

$L \geq$ right-handed trefoil knot (plus trivial components) or




FIGURE 3.5.




FIGURE 3.6.

$L \geq$ right-handed Hopf link (plus trivial components).

Proof. First suppose that the negative crossing N of $\tilde{L}$ is a self-crossing of a component say $\tilde{\ell}$. If $\tilde{L}$ has a mixed crossing then we have $L \geq$ right-handed Hopf link (plus trivial components) by Lemma 2.2. Suppose that $\tilde{\ell}$ has no mixed crossings with other components. The negative crossing N divides $\tilde{\ell}$ into two parts. If the two parts have no common crossings except N , then either we have the right-handed trefoil knot or $\tilde{\ell}$ is almost trivial. Otherwise, choose such a crossing (which is positive) and look at the complementary tangle diagram of the crossing. Note that N is a mixed crossing of that tangle diagram. Then by Lemma 3.1 either we have $\tilde{\ell} \geq$ right-handed trefoil knot or $\tilde{\ell}$ is the diagram $\tilde{T}_2$ of Fig. 1.2.

Next suppose that N is a mixed crossing of the components, say $\tilde{\ell}_1$ and $\tilde{\ell}_2$. If there are components $\tilde{\ell}_i$ and $\tilde{\ell}_j$ with $\tilde{\ell}_i \cap \tilde{\ell}_j \neq \emptyset$ and $\{i, j\} \neq \{1, 2\}$, then we have the right-handed Hopf link. If not, then applying Lemma 2.10 to the complementary tangle projection of N , we have $\tilde{\ell}_1 \cup \tilde{\ell}_2 \geq$ right-handed Hopf link or $\tilde{\ell}_2 \cup \tilde{\ell}_2$ is the diagram $\tilde{T}_3$ of Fig. 1.2. This completes the proof. $\square$

Theorem 1.4, Theorem 1.5, Corollary 1.6 and Corollary 1.7 are immediate corollaries of Theorem 3.2.

4. 2-ALMOST POSITIVE LINKS

We start from some preparatory lemmas that are used to analyze 2-almost positive diagrams. In the following proofs of these lemmas we use sometimes basic facts from Lemma 2.1, Lemma 2.6, Lemma 2.7 etc. from Section 1 without explicitly mentioning it. When we give over/under crossing information to a projection, we divide a projection into some arcs and circles and give over/under crossing information to the self-crossings of these arcs and circles by descending or ascending algorithm without explicitly mentioning it.

Lemma 4.1. *Let $\hat{T}$ be a prime 2-string tangle projection with vertical connection. If $\hat{T}$ is not an underlying projection of any of the tangles in Fig. 4.1, then $\hat{T}$ is one of the projections illustrated in Fig. 4.2*




FIGURE 4.1.




FIGURE 4.2.

Proof. We have that both $\hat{a}$ and $\hat{b}$ are almost trivial by Lemma 2.6. Suppose that there is a rightmost mixed crossing on $\hat{a}$ with its multiplicity greater than one, or a leftmost mixed crossing on $\hat{b}$ with its multiplicity greater than one. Suppose that we have the former case. Then possibly after taking a minor of $\hat{T}$, still denoted by $\hat{T}$ using Lemma 2.1, we find a tear drop disk δ and two rightmost mixed crossings B_i and B_{i+1} on $\partial\delta$ with multiplicity greater than one and $B_i B_{i+1} \subset \delta$. Then we add over/under crossing information to $\hat{T}$ respecting the multiplicity and the position of B_i and B_{i+1} and have the tangle $T(1/2, -2)$ as illustrated in Fig. 4.3.

Suppose that there is a mixed crossing with its depth greater than zero. Then there is a mixed crossing, say B_i on $\hat{a}$ with depth one. Then using Lemma 2.1 we may assume that there is a tear drop disk δ with $B_i \in \partial\delta$. Then by re-choosing B_i if necessary we may further assume that $B_{i+1} \in \partial\delta$ and $B_i B_{i+1} \subset \delta$. We divide $\hat{b}$ into $B_0 B_i$, $B_i B_{i+1}$ and $B_{i+1} B_\infty$ and give over/under crossing information to $\hat{T}$ so that $B_0 B_i^-$ is over everything and $B_{i+1}^+ B_\infty$ is under everything and other crossings are as illustrated in Fig. 4.4 and we have the tangle $T(-4)$.




FIGURE 4.3.




FIGURE 4.4.




FIGURE 4.5.

Therefore we have that $\hat{a}$ is of the form illustrated in Fig. 4.5

Suppose $l > 0$ and the spine $\hat{a}'$ of $\hat{a}$ has mixed crossings. Using Lemma 2.1 we reduce $\hat{a}$ so that $\hat{a}$ bounds just one 1-gon δ . Let V be the vertex of δ .

Case 1 The first mixed crossing B_1 of $\hat{b}$ is on A_0V .

Then we have the tangle $T(-4)$ as illustrated in Fig. 4.6 (a) where B_i denotes the first crossing of $\hat{b}$ at which $\hat{b}$ meets δ .

Case 2. B_1 is on VA_∞ .

Case 2.1. The last crossing of $\hat{b}$ on $B_0B_1 \cup B_1A_\infty$ is on B_0B_1 .

We have the tangle $T(1/2, -2)$ as illustrated in Fig. 4.6 (b).

Case 2.2. The last crossing of $\hat{b}$ on $B_0B_1 \cup B_1A_\infty$ is on B_1A_∞ .

We have the tangle $T(-4)$ as illustrated in Fig. 4.6 (c).

Case 3. B_1 is on $\partial\delta$.

Case 3.1. The first crossing of $\hat{b}$ on $\hat{a}' = A_0V \cup VA_\infty$ is on A_0V .

We have the tangle $T(1/2, -2)$. See Fig. 4.6 (d).

Case 3.2. The first crossing of $\hat{b}$ on $\hat{a}' = A_0V \cup VA_\infty$ is on VA_∞ .

We have the tangle $T(-4)$ as illustrated in Fig. 4.6 (e).

Therefore we have that $\hat{a}'$ has no mixed crossings of $\hat{T}$. Similarly, if $\hat{b}$ is not simple, then $\hat{b}'$ has no mixed crossings of $\hat{T}$. If both $\hat{a}$ and $\hat{b}$ have self-crossings, then we have the tangle $T(1/2, -2)$ by Lemma 2.7. Then we have that at least one of $\hat{a}$ and $\hat{b}$ has no self crossings. Then the result follows from Lemma 2.11 together with Lemma 2.7 and Lemma 2.10. $\square$




FIGURE 4.6.

Lemma 4.2. *Let $\tilde{T}$ be a prime R2-reduced 2-almost positive 2-string tangle diagram with vertical connection. Suppose that $\tilde{T}$ has a negative self-crossing on the left string $\tilde{a}$ and a negative mixed crossing. If $\tilde{T}$ is not greater than or equal to any of the tangles in Fig. 4.7, then $\tilde{T}$ is one of the forms illustrated in Fig. 4.8 where dotted lines have no crossings with the real lines.*




FIGURE 4.7.




FIGURE 4.8.

This lemma is proved by a steady consideration of various cases.

Proof. First we note that the tangles in Fig. 4.7 are closed under horizontal symmetry. In particular $H(T_1) = T_1$. The diagrams in Fig. 4.8 are also closed under horizontal symmetry. We also note here that $\tilde{T}_{4\pm}$ and $\tilde{T}_{5\pm}$, $\tilde{T}_{6\pm}$ and $\tilde{T}_{7\pm}$ are flypes each other respectively. We denote the negative mixed crossing by N and the negative self-crossing by M . As $\tilde{T}$ is not greater than or equal to $T(3_1, 0_1)$, we have by Theorem 3.2 that $\tilde{a}$ is obtained from an R1-augmentation of the diagram $\tilde{T}_1$ or $\tilde{T}_2$ in Fig. 1.2 by deleting a trivial disk pair (D^2, D^1) . As $\tilde{T}$ is not greater than or equal to $T(0_1, 3_1)$, we have by Lemma 2.6 that $\tilde{b}$ is almost trivial.

Case 1. $\tilde{a}$ is obtained from the diagram $\tilde{T}_2$ in Fig. 1.2.

Suppose $\tilde{a}$ is obtained from an R1-augmentation $\tilde{S}_2$ of $\tilde{T}_2$ by deleting a trivial disk pair.

Suppose that a component of $(\tilde{S}_2 - \tilde{T}_2) \setminus \text{int} D^2$ is incident to ∂D^2 and has a mixed crossing of $\tilde{T}$. Since $\tilde{T}$ is R2-reduced we can apply Lemma 3.1 after erasing the negative self-crossing M using Lemma 2.2 and have the tangle $T(-2)$. See Fig. 4.9.




FIGURE 4.9.

By similar arguments based on Lemma 2.2 and Lemma 3.1 we may suppose that $\tilde{a}$ is an R1-augmentation of one of the diagrams illustrated in Fig. 4.10 and their horizontal symmetries, and that both A_0P_+ and P_+A_∞ have no mixed crossings for $p > 0$.




FIGURE 4.10.

We remark that as $\tilde{T}$ is prime R2-reduced, both $s(\tilde{a}_1, Q_+)$ and the 2-gon UVU of $\tilde{a}_{2\pm}$ have mixed crossings. We will show that in Case 1 $\tilde{T} \geq T(-2), T_1, T_3, T_4, T_5$ or T_6 .

As A_0P_+ and P_+A_∞ have no mixed crossings we may suppose, by taking a flype of $\tilde{T}$, still denoted by $\tilde{T}$ if necessary that $p = 0$.

Case 1.1. $\tilde{a}$ is an R1-augmentation of the diagram $\tilde{a}_1$ of Fig. 4.10 with $p = 0$ and q odd. Then we have $U = M$. See Fig. 4.11.

Suppose that N is on $s(\tilde{a}, M)$. If $r(\tilde{a}, M)$ have mixed crossings then we have the tangle $T(-2)$ by Lemma 2.2 and Lemma 2.7. If not then we have $q \geq 3$ by the R2-reducibility. Note that by Lemma 2.2 we have that $r(\tilde{T}, V)$ is a minor of $\tilde{T}$. Then by applying Lemma 3.1 to $r(\tilde{T}, V)$ we have the tangle $T(-2), T_3$ or T_4 . See Fig. 4.12.

Therefore we may suppose that N is on $r(\tilde{a}, M)$. Since $r(\tilde{a}, M)$ is a minor of $\tilde{T}$ by Lemma 2.2 we may assume by Lemma 3.1 that $r(\tilde{T}, M)$ is an R1




FIGURE 4.11.




FIGURE 4.12.

augmentation of one of the diagrams $\tilde{T}_{1+}, \tilde{T}_{1-}, \tilde{T}_{2+}$ and $\tilde{T}_{2-}$ in Fig. 3.2. We will show that $\tilde{T}$ is greater than or equal to $T(-2)$ or T_1 in all cases as follows.

Case 1.1.1. The first crossing of $\tilde{b}$ with $r(\tilde{a}, M)$ is N and is on A_0M .

We have $\tilde{T} \geq T(-2)$ as illustrated in Fig. 4.13. Here Fig. 4.13 (a) describes the case that N is not the first mixed crossing of $\tilde{b}$. The case N is the first mixed crossing is illustrated in other figures. Because $\tilde{T}$ is prime and R2-reduced, we have one of the situations in Fig. 4.13. Note that there are at most two dotted lines in each figure so that we can trivialize them by adding over/under crossing information as one is over everything and the other is under everything. We also note that the crossings may be on the R1 residual of $\tilde{a}$ with respect to $\tilde{a}_1$ of Fig. 4.10. That case is not illustrated in Fig. 4.13 for the simplicity. However it reduces to the case that the crossings are on the core. See Fig. 4.14 how to reduce it. In Fig. 4.14 (a) the dotted part is supposed to be given over/under crossing information so that it is under everything. In Fig. 4.14 (b), (c), (d), (e) the dotted line

bounded by two marked crossings are contained in the tear drop disk. Then we have the same result as (a) keeping the conditions that another dotted line is given over/under crossing information so that it is under everything.




FIGURE 4.13.

Case 1.1.2. The first crossing of $\tilde{b}$ with $r(\tilde{a}, M)$ is N and on MQ_+ .

By applying Lemma 3.1 to $r(\tilde{T}, M)$ we may suppose that $\tilde{a}$ is over $\tilde{b}$ at N . Then we may suppose that $r(\tilde{a}, Q_+) \cap \tilde{b} = \emptyset$, and $r(\tilde{a}, V) \cap \tilde{b} = \emptyset$ otherwise we have $\tilde{T} \geq T(-2)$ by Lemma 2.2 and Lemma 2.7. Then we have $\tilde{T} \geq T(-2)$ as illustrated in Fig. 4.15

Case 1.1.3. The first crossing of $\tilde{b}$ with $r(\tilde{a}, M)$ is N and on $Q_+Q_-A_\infty$.

If the first mixed crossing of $\tilde{b}$ is not N then we have $T(-2)$ as illustrated in Fig. 4.16 (a). If N is the first mixed crossing of $\tilde{b}$ then we have the situation illustrated in Fig. 4.16 (b) and then also have $T(-2)$.

Case 1.1.4. The last crossing of $\tilde{b}$ with $r(\tilde{a}, M)$ is N and on VA_∞ . See Fig. 4.17.

Case 1.1.5. The last crossing of $\tilde{b}$ with $r(\tilde{a}, M)$ is N and on $A_0M \cup MQ_+Q_-V$. See Fig. 4.18.

Case 1.2. $\tilde{a}$ is an R1-augmentation of the diagram $\tilde{a}_1$ of Fig. 4.10 with $p = 0$ and q even, $q \geq 2$. Then we have $V = M$. See Fig. 4.19.

Suppose $s(\tilde{a}, U) \cup s(\tilde{a}, M)$ has no mixed crossings. Then using Lemma 3.1 to $r(r(\tilde{T}, U), M)$ we have the tangle $T(-2)$ as illustrated in Fig. 4.20.

Next suppose $s(\tilde{a}, U)$ has two positive crossings. Then we have $\tilde{T} \geq r(\tilde{T}, M) \geq T(-2)$ by Lemma 2.2 and Lemma 3.1. Thus we may suppose that $s(\tilde{a}, U)$ does not have two positive crossings. If N is on $s(\tilde{a}, U) \cup s(\tilde{a}, M)$ and $r(r(\tilde{a}, U), M)$ has mixed crossings then we also have $T(-2)$.




FIGURE 4.14.




FIGURE 4.15.




FIGURE 4.16.




FIGURE 4.17.




FIGURE 4.18.




FIGURE 4.19.

If $r(r(\tilde{a}, U), M)$ has no mixed crossings then because $\tilde{T}$ is R2-reduced, we have $s(\tilde{a}, U) \cup s(\tilde{a}, M)$ has four mixed crossings. Then $s(\tilde{a}, M)$ has two positive mixed crossings. If N is on $r(r(\tilde{a}, U), M)$ then by the primeness of $\tilde{T}$ we have that $s(\tilde{a}, M)$ has two positive mixed crossings. Therefore we




FIGURE 4.20.

may suppose that $s(\tilde{a}, M)$ has two positive mixed crossings. Then we may suppose that they are successive on $\tilde{b}$. Namely the arc of $\tilde{b}$ bounded by them are contained in a region bounded by $s(\tilde{a}, M)$. We may also assume by taking a minor of $\tilde{T}$, still denoted by $\tilde{T}$ using Lemma 2.2 that these two positive mixed crossings considered on $r(\tilde{a}, U)$ are rightmost. Note that we may also suppose that either $s(\tilde{a}, U)$ has no mixed crossings or it has just two mixed crossings, one is positive the other is N . Then we have $T(-2)$ or T_1 by “over and under” or “under and over” technique used in the proof of Lemma 3.1 in the cases illustrated in Fig. 4.21 and Fig. 4.22.

The rest is the case of Fig. 4.23.

Note that in this case $s(\tilde{a}, U)$ has two mixed crossings and therefore N is on it. Then either we can apply Lemma 2.9 for a sub-tangle diagram of $r(\tilde{T}, U)$ to have the tangle $T(-2)$ as illustrated in Fig. 4.24 or we have the tangle $T(-2)$ or T_3 as illustrated in Fig. 4.25.

Case 1.3. $\tilde{a}$ is an R1-augmentation of the diagram $\tilde{a}_{2\pm}$ in Fig. 4.10 with $p = 0$ or its horizontal symmetry. See Fig. 4.26.

We may suppose without loss of generality that $q \geq r$ by horizontal symmetry. We will show that we have the tangle $T(-2)$, T_1 or T_5 in this case. Suppose that $Q_-Q_+ \cup Q_+Q_- \cup S_-S_+ \cup S_+S_-$ has mixed crossings. Suppose for example that there are two successive mixed crossings P_1 and P_2 . If one of them is N then by Lemma 2.2 we have $\tilde{T} \geq r(\tilde{T}, S_+)$ and taking the fact that the 2-gon UVU has mixed crossings then because $\tilde{T}$ is R2-reduced, we have $r(\tilde{T}, S_+) \geq T(-2)$ by Lemma 3.1. Suppose that both P_1 and P_2 are positive crossings. Then by “over and under technique” we have $\tilde{T} \geq T(-2)$. Thus we have that $Q_-Q_+ \cup Q_+Q_- \cup S_-S_+ \cup S_+S_-$ has no mixed crossings. Therefore we have that any $\tilde{T}$ is greater than or equal to a tangle, still denoted by $\tilde{T}$ of the following:

$q = 0$ and $s = 2$, or $q = 0$ and $s = 1$, or $q = s = 1$.

Case 1.3.1 $q = 0$ and $s = 2$.

For the diagram $\tilde{a}_{2-}$ of Fig. 4.26, we have $U = M$ and have the tangle $T(-2)$ from $r(\tilde{T}, M)$ by Lemma 3.1. Therefore we consider the diagram of Fig. 4.27.

If N is on $s(\tilde{a}, M)$ then we have $T(-2)$. If N is not on $s(\tilde{a}, M)$ but on $s(\tilde{a}, S_+)$ and $\tilde{a}$ is over $\tilde{b}$ at N then we have $T(-2)$ by applying Lemma 2.7 to $r(\tilde{T}, S_-)$. Thus we have that $\tilde{a}$ is under $\tilde{b}$ at N . Then applying Lemma 3.1 to $r(\tilde{T}, M)$ we have the result as illustrated in Fig. 4.28. Note that Fig. 4.28 (a), (b) and (c) do not occur since $\tilde{T}$ is R2-reduced. Note that Fig. 4.28 (b) and (c) are the cases that $r(\tilde{T}, M)$ has just two mixed crossings.




FIGURE 4.21.

Suppose N is on otherwise. Then there are at least two positive mixed crossings on $s(\tilde{a}, S_+)$. Then we have the result by “over and under technique” as illustrated in Fig. 4.29 or Fig. 4.30. Note that in the case illustrated in Fig. 4.29 (a) we have that $\tilde{T} \geq r(\tilde{T}, S_-) \geq T(-2)$ and the case illustrated in Fig. 4.29 (b) reduces to other cases. In Fig. 4.30 we describe the situation that $s(\tilde{a}, S_+) \setminus s(\tilde{a}, M)$ has no mixed crossings. Then, because $\tilde{T}$ is R2-reduced, we have that the middle dotted line of $\tilde{b}$ illustrated in Fig. 4.30 (a) and (b) must intersect MU . Then applying Lemma 3.1




FIGURE 4.22.




FIGURE 4.23.

to $r(\tilde{T}, S_-)$ we have the situation illustrated in Fig. 4.30 (c) or (d) and we have that $\tilde{T}$ is greater than or equal to $T(-2)$.

Case 1.3.2. $q = 0$ and $s = 1$.

First we consider the case that $\tilde{a} = \tilde{a}_{2+}$ in Fig. 4.26. Then we have $M = U$. See Fig. 4.31.

Suppose that N is on $s(\tilde{a}, M)$. Since $\tilde{T}$ is R2-reduced we have that $r(\tilde{a}, M)$ has mixed crossings. Then by Lemma 2.2 and Lemma 2.7 we have $T(-2)$. Thus we have that N is not on $s(\tilde{a}, M)$. Suppose that N is on MS in $s(\tilde{a}, S)$. If $\tilde{a}$ is over $\tilde{b}$ at N then we have $\tilde{T} \geq r(\tilde{T}, S)$ by Lemma 2.2 and we have $r(\tilde{T}, S) \geq T(-2)$ by Lemma 2.7. Thus we have that $\tilde{a}$ is




FIGURE 4.24.




FIGURE 4.25.




FIGURE 4.26.




FIGURE 4.27.




FIGURE 4.28.

under $\tilde{b}$ at N . Then by applying Lemma 3.1 to $r(\tilde{T}, M)$ we have that N is the last mixed crossing of $\tilde{b}$ with $r(\tilde{T}, M)$. Therefore we have the situation illustrated in Fig. 4.32 and have the tangle $T(-2)$.

Thus we have that N is not on $s(\tilde{a}, S)$. Then we have that there are two successive positive mixed crossings on $s(\tilde{a}, S)$. If they are not on VM of $s(\tilde{a}, S)$ then we have $T(-2)$ by “over and under technique”. We also have $T(-2)$ in the cases illustrated in Fig. 4.33 (a) and (b). For the case illustrated in Fig. 4.33 (c) we do not have the situation illustrated in Fig. 4.33 (d) because in this situation the second dotted line has a mixed crossing with $r(\tilde{a}, S)$ and then we have $r(\tilde{a}, S) \geq T(-2)$ by Lemma 3.1. Thus we have that the case illustrated in Fig. 4.33 (c) reduces to other cases. In the case illustrated in Fig. 4.33 (e) we have by taking the position of N on $r(\tilde{a}, M)$ into account using Lemma 3.1 we have that the first dotted line of $\tilde{b}$ can be under everything, the third dotted line can be over everything, and the second dotted line can be over everything except the third dotted line. Thus we have the tangle $T(-2)$ as illustrated. The final case is the case that the two successive positive mixed crossings are on VM of $s(\tilde{a}, S)$. Then because $\tilde{T}$ is R2-reduced, we have that the arc of $\tilde{b}$ bounded by them




FIGURE 4.29.




FIGURE 4.30.

intersects VM of $r(\tilde{a}, M)$. Then by applying Lemma 3.1 to $r(\tilde{T}, M)$ and $r(\tilde{T}, S)$, and because $\tilde{T}$ is R2-reduced, we have the situation illustrated in Fig. 4.33 (f) and then have the tangle $T(-2)$ as illustrated.

Next we consider the case that $\tilde{a} = \tilde{a}_{2-}$ in Fig. 4.26. Then we have $M = V$. See Fig. 4.34.




FIGURE 4.31.




FIGURE 4.32.

First suppose that N is on $s(\tilde{a}, M)$. If $r(\tilde{a}, M)$ has mixed crossings then we have $T(-2)$. Therefore we have $r(\tilde{a}, M)$ has no mixed crossings. Suppose that N is on MU of $s(\tilde{a}, M)$. Then after changing some mixed crossings on MU of $s(\tilde{a}, M)$ except N and deforming MU of $s(\tilde{a}, M)$ away from $s(\tilde{a}, S)$ we have the tangle $T(-2)$ by the use of Lemma 2.7. Therefore we may suppose that N is on SM . Then we may suppose that $\tilde{a}$ is over $\tilde{b}$ at N , otherwise we can apply Lemma 2.2 to $r(\tilde{T}, S)$. Let P be the first mixed crossing of $\tilde{b}$ with MU of $s(\tilde{a}, M)$. Then we have the tangle T_1 as illustrated in Fig. 4.35 (a), or we have the situation illustrated in Fig. 4.35 (b). Then we give over/under crossing information so that the third dotted line is under everything, the first dotted line is under everything except the third dotted line, and the second dotted line is over everything. Note that we can first deform the third dotted line such that it does not intersect the first dotted line as illustrated in Fig. 4.35 (c) and we have the tangle T_1 .

Thus we have that N is not on $s(\tilde{a}, M)$. Now suppose that N is on $s(\tilde{a}, S)$. Then we have one of the situations illustrated in Fig. 4.36 and we have $T(-2)$ or T_1 . Note that in the situation illustrated in Fig. 4.36 (c) there are just two mixed crossings of $r(\tilde{T}, M)$.

Thus we have that N is not on $s(\tilde{a}, M) \cup s(\tilde{a}, S)$. Then there are two successive positive mixed crossings on $s(\tilde{a}, S)$. If they are not on MU then we have $T(-2)$ by “over and under technique”. If they are as illustrated in Fig. 4.37 (a), (b), (c) or (d) then we have $T(-2)$ or T_1 . Then we assume that there are no such situations. In the situation illustrated in Fig. 4.38 (a) we either have T_1 , or if the third dotted line do not intersects SA_∞




FIGURE 4.33.




FIGURE 4.34.




FIGURE 4.35.




FIGURE 4.36.

then we deform A_0MUS first and then we have $T(-2)$ as illustrated. If the third dotted line intersects SA_∞ then we have the situation illustrated in Fig. 4.38 (b) and we have $T(-2)$. In the case illustrated in Fig. 4.38 (c) we apply Lemma 3.1 to $r(\tilde{T}, M)$ and we have $T(-2)$.

Case 1.3.3. $q = s = 1$.

First we consider the case that $\tilde{a} = \tilde{a}_{2+}$ in Fig. 4.26. Then we have $M = V$. By Lemma 2.2 we have $\tilde{T} \geq r(\tilde{T}, M)$. Note that, because $\tilde{T}$ is R2-reduced, the 2-gon UMU has mixed crossings. Then by Lemma 3.1 we have that $r(\tilde{T}, M) \geq T(-2)$.

Next we consider the case that $\tilde{a} = \tilde{a}_{2-}$ in Fig. 4.26. Then we have $M = U$. See Fig. 4.39.

Note that, because $\tilde{T}$ is R2-reduced, the 2-gon MVM has mixed crossings. Therefore we have that both $s(\tilde{a}, Q)$ and $s(\tilde{a}, S)$ have mixed crossings. Suppose that N is on $r(r(\tilde{a}, Q), S)$. By Lemma 2.2 we have $\tilde{T} \geq r(\tilde{T}, Q)$ and $\tilde{T} \geq r(\tilde{T}, S)$. Then by Lemma 3.1 we have $r(\tilde{T}, Q) \geq T(-2)$. Therefore we have that N is not on $r(r(\tilde{a}, Q), S)$. Then up to horizontal symmetry we may suppose that N is on $s(\tilde{a}, S)$. By Lemma 2.2 we have $\tilde{T} \geq r(\tilde{T}, Q)$. If $\tilde{a}$ is under $\tilde{b}$ at N then by Lemma 2.2 we have $\tilde{T} \geq r(\tilde{T}, S)$. Then since $r(\tilde{T}, S)$ is a positive diagram we have $r(\tilde{T}, S) \geq T(-2)$ by Lemma 2.7. Therefore we have that $\tilde{a}$ is over $\tilde{b}$ at N . We have by Lemma 3.1 that $r(\tilde{T}, Q)$ is an R1-augmentation of the diagram $\tilde{T}_{1+}$ in Fig. 3.2. Now we consider the position of two successive positive mixed crossings on $s(\tilde{a}, Q)$. If they are not on MV then we have $T(-2)$ by “over and under technique”. Suppose that one of them is on QM and the other is on MV . Since $s(\tilde{a}, A)$ has just two mixed crossings we have that this case reduces to the previous case. Suppose that one of them is on MV and the other is on VQ . Then we have $T(-2)$ or T_5 as illustrated in Fig. 4.40 (a) and (b). Finally suppose that no former cases occur and both of them are on MV . Then we consider the position of the first mixed crossing of $\tilde{b}$ with $s(\tilde{a}, Q)$, the last mixed crossing of $\tilde{b}$ with $s(\tilde{a}, Q)$ and the two mixed crossings of $\tilde{b}$ with $s(\tilde{a}, S)$. Then we have $T(-2)$ or T_1 as illustrated in Fig. 4.40 (c).




FIGURE 4.37.




FIGURE 4.38.




FIGURE 4.39.




FIGURE 4.40.

Case 2. $\tilde{a}$ is almost trivial.

By Lemma 3.1 we may suppose that $r(\tilde{T}, M)$ is up to horizontal symmetry, an R1-augmentation of one of the diagrams in Fig. 4.41. Note that because $\tilde{T}$ is R2-reduced, $r(\tilde{a}, M)$ has no R1-residuals.

Case 2.1. Suppose that $B_0 Q_+ \cup Q_+ B_\infty$ has mixed crossings. Then by applying Lemma 2.7 to $r(\tilde{T}, Q_+)$ we have the tangle $T(-2)$. Therefore we have that $B_0 Q_+ \cup Q_+ B_\infty$ has no mixed crossings. Then by flyping we may suppose that $q = 0$.




FIGURE 4.41.

In the following we consider the case $p \leq 1$. By that argument it is easily seen that in the case $p \geq 2$ we have the tangle $T(-2)$.

First suppose $p = 1$. We consider the position of M on $\tilde{a}$ and where $\tilde{b}$ $s(\tilde{a}, M)$ has intersection with $\tilde{b}$. A dotted arrow in Fig. 4.42 implies that M is on its initial point and $s(\tilde{a}, M)$ has intersection with the part of $\tilde{b}$ where the terminal point is there. For example Arrow (1) describes the case that M is on UP and $s(\tilde{a}, M)$ has intersection with UN of $\tilde{b}$. We denote this case by Case (1). Because $\tilde{T}$ is R2-reduced, there are only five possible cases illustrated in Fig. 4.42. Note that Case (3) and Case (5) may occur simultaneously. In Cases (1), (2) and (3) we have the tangle $T(-2)$ by Lemma 2.7 as illustrated in Fig. 4.43. Thus we have that these cases do not happen. In Case (4) we have $T(-2)$ by Lemma 2.9 unless $\tilde{T}$ is a diagram $\tilde{T}_{4+}$ in Fig. 4.8 as illustrated in Fig. 4.43. In Case (5) we have $T(-2)$ by Lemma 2.9 unless $\tilde{T}$ is a diagram $\tilde{T}_{6+}$ in Fig. 4.8 as illustrated in Fig. 4.43.




FIGURE 4.42.

Next suppose $p = 0$. Then we have the cases illustrated in Fig. 4.44. We have $T(-2)$ in Cases (1) and (2). Cases (3), (4) and (5) correspond the diagrams $\tilde{T}_{3+}$, $\tilde{T}_{2++}$ and $\tilde{T}_{2-+}$ in Fig. 4.8 respectively.

Case 2.2. In the cases illustrated in Fig. 4.44 we have the tangle $T(-2)$ or T_2 as illustrated in Fig. 4.45. Here we denote by Case (i)' a subcase of Case (i). The proof for Case (i)' is entirely analogous to that of Case (i) after removing extra crossings. Other cases are illustrated in Fig. 4.46. Then we have in Cases (6), (7), (8), (9) and (10) the diagram $\tilde{T}_{2-+}$ in Fig. 4.8 and in Cases (11), (12), (13), (14) and (15) the diagram $\tilde{T}_{2++}$ in Fig. 4.8.

Case 2.3. In this case we have the diagram $\tilde{T}_1$ in Fig. 4.8.




FIGURE 4.43.




FIGURE 4.44.

This completes the proof of Lemma 4.2. □

Lemma 4.3. *Let $\hat{T}$ be a 2-string tangle projection with vertical connection. Suppose that $\hat{T}$ has no self-crossings. If $\hat{T}$ is not a projection of the tangle $R(T(3, 1))$ illustrated in Fig. 4.48 then $\hat{T} = \hat{T}(2n)$ (see Fig. 2.14) for some non-negative integer n .*

Proof. The proof here is given by contradiction. Suppose that $\hat{T}$ is a 2-string tangle projection with vertical connection with $2n$ mixed crossings and no self-crossings that is not equal to $\hat{T}(2n)$ such that it is not a projection of the tangle $R(T(3, 1))$. Suppose that n is smallest among such tangle projections. It is clear that $\hat{T}$ has a 2-gon as $\hat{T}_1$ illustrated in Fig. 2.17. Let $\hat{T}'$ be a tangle projection obtained from $\hat{T}$ by replacing the 2-gon by a pair of parallel arcs as $\hat{T}_2$ illustrated in Fig. 2.17. Then by Lemma 2.4 we have that $\hat{T}' = \hat{T}(2(n - 1))$. However it is easy to check that all




FIGURE 4.45.




FIGURE 4.46. Case 2.2 of the proof of Lemma 4.2

tangle projections without mixed crossings obtained from $\hat{T}(2(n-1))$ by replacing a pair of parallel arcs by a 2-gon except $\hat{T}(2n)$ is a projection of $R(T(3,1))$. See Fig. 4.47. This completes the proof. $\square$

Lemma 4.4. *Let $\hat{T}$ be a reduced 2-string tangle projection with vertical connection. If $\hat{T}$ is not a projection of any of the tangles in Fig. 4.48 then $\hat{T}$ is one of the projections illustrated in Fig. 4.49*




FIGURE 4.47.




FIGURE 4.48.

Proof. We have that both $\hat{a}$ and $\hat{b}$ are almost trivial.

First suppose that $\hat{a}$ has self-crossings. If there is a self-crossing of $\hat{a}$ which is not rightmost then we have $R(T(3, 1))$ using Lemma 2.1 as illustrated in Fig. 4.50.

Therefore all mixed crossings on $\hat{a}$ are rightmost. Let B_{2i+1} be a mixed crossing with maximal multiplicity among all mixed crossings on $\hat{a}$. Then we have that B_{2i+2} also has the maximal multiplicity. We consider the position of successive mixed




FIGURE 4.49.




FIGURE 4.50.

crossings B_{2i} , B_{2i+1} and B_{2i+2} , or B_{2i+1} , B_{2i+2} and B_{2i+3} . If they are positioned as illustrated in Fig. 4.51, where only the positions of B_{2i} , B_{2i+1} and B_{2i+2} are illustrated, then we have the tangle $R(T(3, 1))$. Up to horizontal symmetry we have the tangle $R(T(3, 1))$ for the positions of B_{2i+1} , B_{2i+2} and B_{2i+3} corresponding to that of B_{2i} , B_{2i+1} illustrated in Fig. 4.51. Note that in Fig. 4.51 the position of B_{2i} is not precisely described. Namely the multiplicity of B_{2i} on $\hat{a}$ may be greater than that illustrated in Fig. 4.51. However by the argument described in Fig. 4.14 we have the same result for these cases.




FIGURE 4.51.

Note that all cases that the multiplicity $m(B_{2i+1})$ is odd are illustrated in Fig. 4.51 (a), (b), (c), (d) and (e). Suppose that B_{2j} , B_{2i+1} and B_{2i+2} are positioned as illustrated in Fig. 4.52 (a) where $j \leq i$ and B_k are related to B_{2i+1} on $\hat{a}$ for $2j < k \leq 2i$. Let P be the root of B_{2i+1} on $\hat{a}$. Then we have that $B_{2j}B_{2i+1}$ and $A_0P \cup PA_\infty$ intersect only at B_{2j} . Therefore after deforming B_0B_{2j} as illustrated in Fig. 4.52 (a) we have the tangle $R(T(3, 1))$. Suppose that B_{2i} and B_{2i+1} are positioned as illustrated in Fig. 4.52 (b). Then either we have the case illustrated in Fig. 4.52 (a) or the case that is the horizontal symmetry of the case illustrated

in Fig. 4.51 (f) or (i), or we have the case illustrated in Fig. 4.52 (c). In any case we have $R(T(3, 1))$.




FIGURE 4.52.

Therefore we have that all mixed crossings are related on $\hat{a}$. Similarly we have that all mixed crossings are related on $\hat{b}$. Therefore we have the situation that there is a subangle projection $\hat{T}_0$ of $\hat{T}$ as illustrated in Fig. 4.53 (a) where m and n are non-negative integers such that $\hat{T}_0$ has no self-crossings.

First suppose that $m + n$ is odd. If $\hat{T}_0$ is a projection of the tangle $T(4)$ then we have $R(T(3, 1))$ as illustrated in Fig. 4.53 (b). Then we have the conclusion by applying Lemma 2.11 to $\hat{T}_0$.

Next suppose that $m + n$ is even. If $\hat{T}_0$ is a projection of the tangle $R(T(3, 1))$ then we clearly have that $\hat{T}$ is also a projection of $R(T(3, 1))$. Then we have the conclusion by applying Lemma 4.3 to $\hat{T}_0$. This completes the proof. $\square$




FIGURE 4.53.

Lemma 4.5. *Let $\hat{T}$ be a 2-string tangle projection with X-connection. Suppose that $\hat{T}$ has no self-crossings. Suppose that $\hat{T}$ is not a projection of the tangle $T(1/2, -3)$ illustrated in Fig. 4.56. Then there are odd numbers $n_1, n_2, \dots, n_k$ such that $\hat{T} = R(\hat{T}(n_1, n_2, \dots, n_k))$ (Fig. 4.54).*




FIGURE 4.54.

Proof. The proof here is given by contradiction. Suppose that $\hat{T}$ is a 2-string tangle projection with X-connection with n mixed crossings and no self-crossings that is not equal to any $R(\hat{T}(n_1, n_2, \dots, n_k))$ such that it is not a projection of the tangle $T(1/2, -3)$. Suppose that n is smallest among such tangle projections. It is clear that $\hat{T}$ has a 2-gon as $\hat{T}_1$ illustrated in Fig. 2.17. Let $\hat{T}'$ be a tangle projection obtained from $\hat{T}$ by replacing the 2-gon by a pair of parallel arcs as $\hat{T}_2$ illustrated in Fig. 2.17. Then by Lemma 2.4 we have that $\hat{T}' = R(\hat{T}(n_1, n_2, \dots, n_k))$ for some odd numbers $n_1, n_2, \dots, n_k$. However it is easy to check that a tangle projection without mixed crossings obtained from $R(\hat{T}(n_1, n_2, \dots, n_k))$ by replacing a pair of parallel arcs by a 2-gon is either a projection of $T(1/2, -3)$, or equal to $R(\hat{T}(n'_1, n'_2, \dots, n'_{k'}))$ for some odd numbers $n'_1, n'_2, \dots, n'_{k'}$. See Fig. 4.55. In Fig. 4.55 some typical cases are illustrated. Any other case is essentially the same as one of the typical cases. In Fig. 4.55 (a) some typical cases that will produce some $R(\hat{T}(n'_1, n'_2, \dots, n'_{k'}))$ are illustrated. This completes the proof. $\square$

Lemma 4.6. *Let $\hat{T}$ be a reduced 2-string projection with X-connection which is not a projection of any of the tangles in Fig. 4.56. Then $\hat{T}$ is (a flype of) (the vertical and/or horizontal symmetry of) one of the tangle projections in Fig. 4.57.*

Proof. By Lemma 4.5 and by the fact that $\hat{T}$ is not a projection of $T(1/2, -3)$, $T(-5)$ or $R(T(3, 1, 3))$ we have that the core $\hat{T}' = \hat{a}' \cup \hat{b}'$ is (a flype of) a projection in Fig. 4.58.

Case 1. If $\hat{T}$ is not equal to (a flype of) $\hat{T}_2$ then $\hat{T}$ has self-crossing. Then using Lemma 2.7 we have the tangle $T(-5)$ as typically illustrated in Fig. 4.59 and we have the conclusion. In Fig. 4.59 a dotted line expresses that there is a self-crossing, say P of $\hat{T}$ on the core $\hat{T}'$ such that P is on one end of the dotted line and $s(\hat{T}, P)$ has an intersection with a part of $\hat{T}$ where the other end is on. Up to symmetry and flying Fig. 4.59 illustrates all cases.




FIGURE 4.55.

Case 2.1 $\hat{T}'$ has three or more mixed crossings.

If $\hat{T}$ is not equal to (a flype of) $\hat{T}_3$ then using Lemma 2.7 and Lemma 2.9 we have the tangle $T(1/2, -3)$, $R(T(3, 1, 3))$ or T_1 as illustrated in Fig. 4.60.

Case 2.2. $\hat{T}'$ has just one mixed crossing.

First suppose that there is a self-crossing P on the core $\hat{T}'$ such that $r(\hat{T}, P)$ has just one mixed crossing. Then by we apply Lemma 4.1 to a certain subtangle projection of $\hat{T}$ and have either $\hat{T}$ is a projection of




FIGURE 4.56.




FIGURE 4.57.




FIGURE 4.58.




FIGURE 4.59.




FIGURE 4.60.

$T(-5)$, $T(1/2, -3)$, $R(T(3, 1, 3))$ or T_1 , or $\hat{T}$ is equal to (a flype of) $\hat{T}_4$, $\hat{T}_5$ or $\hat{T}_6$. See Fig. 4.61.

Next suppose that for any self-crossing P on the core $\hat{T}' r(\hat{T}, P)$ has three or more mixed crossings. Then we take self-crossings $P_1, P_2, \dots, P_n$ on the




FIGURE 4.61.

core $\hat{T}'$ such that $r(r(\cdots r(\hat{T}, P_1), P_2), \cdots, P_{n-1})$ has three or more mixed crossings and $r(r(\cdots r(\hat{T}, P_1), P_2), \cdots, P_n)$ has just one mixed crossing. Set $\hat{S}_0 = r(r(\cdots r(\hat{T}, P_1), P_2), \cdots, P_{n-2})$ and $\hat{S} = r(r(\cdots r(\hat{T}, P_1), P_2), \cdots, P_{n-1})$. Then we have that $\hat{S} = r(\hat{S}_0, P_{n-1})$ is equal to (a flype of) $\hat{T}_4$, $\hat{T}_5$ or $\hat{T}_6$. Then by applying Lemma 2.7 we have the tangle $T(-5)$, $T(1/2, -3)$, or T_1 as typically illustrated in Fig. 4.62, 4.63 and 4.64. Note that in Fig. 4.62, 4.63 and 4.64 a dotted line express the situation that P_{n-1} is at one end and $s(\hat{S}_0, P_{n-1})$ has intersection with the part of $\hat{S}$ where the other end is on. All typical cases up to flyping are illustrated in Fig. 4.62, 4.63 and 4.64 and other cases are entirely analogous. This completes the proof.

□




FIGURE 4.62.




FIGURE 4.63.

Lemma 4.7. *Let $\tilde{T}$ be an almost positive 2-string tangle diagram with X -connection. Suppose that $\tilde{T}$ has no self-crossings and $\tilde{T}$ is not greater than or equal to the tangle $T(-3)$ nor $T(1/3)$ (Fig. 4.65). Then the underlying projection $\hat{T}$ of $\tilde{T}$ is equal to $\hat{T}(1)$, $\hat{T}(3)$ or $\hat{T}(1/3)$.*




FIGURE 4.64.

Proof. We give the proof by the contradiction. Suppose that there is an almost positive 2-string tangle diagram $\tilde{T}$ with X-connection without self-crossings that is not greater than or equal to the tangle $T(-3)$ nor $T(1/3)$ that contains five or more mixed crossings. We take such $\tilde{T}$ with minimal mixed crossings. It is clear that a tangle with X-connection without self-crossings with three or more mixed crossings has a 2-gon. Thus we have that $\tilde{T}$ has a 2-gon. If one of the two crossings of the 2-gon is the negative crossing then we have a positive diagram $\tilde{T}_0$ from $\tilde{T}$ by applying the second Reidemeister move. Then $\tilde{T}_0$ represents the same tangle as $\tilde{T}$ and has three or more mixed crossings. Then by Lemma 2.10 we have that $\tilde{T}_0$ is greater than or equal to $T(-3)$ or $T(1/3)$. Thus we have that this case does not happen. Therefore we have that the crossings of the 2-gon are both positive. Let $\tilde{T}_1$ be a diagram obtained from $\tilde{T}$ by replacing that 2-gon by a pair of parallel arcs. It is clear that $\tilde{T}$ is greater than or equal to $\tilde{T}_1$. Therefore we have that $\tilde{T}_1$ is not greater than or equal to the tangle $T(-3)$ nor $T(1/3)$. By the minimality of $\tilde{T}$ we have that $\tilde{T}_1$ has just three mixed crossings. Therefore we have that the underlying projection of $\tilde{T}_1$ is $\hat{T}(3)$ or $\hat{T}(1/3)$ and by considering the position of the negative crossing we have that $\tilde{T}_1$ is one of the six tangle diagrams whose underlying projection is $\hat{T}(3)$ or $\hat{T}(1/3)$. But then it is easy to check that any tangle diagram obtained from $\tilde{T}_1$ by replacing a pair of parallel arcs by a 2-gon with positive crossings is a diagram of $T(-3)$ or $T(1/3)$. This is a contradiction. $\square$

Lemma 4.8. *Let $\tilde{T}$ be a prime R2-reduced almost positive 2-string tangle diagram with X-connection. Suppose that the negative crossing of $\tilde{T}$ is a mixed crossing and $\tilde{T}$ is not greater than or equal to any of the tangles in Fig. 4.65. Then $\tilde{T}$ is one of the diagrams illustrated in Fig. 4.66.*




FIGURE 4.65.




FIGURE 4.66.

Proof. We consider the following cases.

Case 1. There is a self-crossing P such that $r(\tilde{T}, P)$ is a positive diagram. Suppose that there are three or more mixed crossings of $r(\tilde{T}, P)$. Then by Lemma 2.10 we have $T(1/3)$ or $T(-3)$. Thus we have that $r(\tilde{T}, P)$ has just one mixed crossing. Then by Lemma 3.1 we have the tangle $T(-3)$ or $T(1/3)$ as illustrated in Fig. 4.67.

Case 2. $\tilde{T}$ has no self-crossings.

In this case we have $\tilde{T}$ is equal to $\tilde{T}_1$ by Lemma 4.7.

Case 3. Otherwise.

By Case 2 we have that the spine $\tilde{T}'$ of $\tilde{T}$ is equal to $\tilde{T}_1$. Then by applying Lemma 2.9 or Lemma 2.7 as illustrated in Fig. 4.68 we have the result. This completes the proof.




FIGURE 4.67.




FIGURE 4.68.

□

Lemma 4.9. *Let $\tilde{L} = \tilde{\ell}_1 \cup \tilde{\ell}_2$ be a 2-almost positive link diagram of a 2-component link L such that the negative crossings are mixed crossings. Suppose that $\tilde{L}$ has four or more mixed crossings. Then L is greater than or equal to the 2-component trivial link.*

Proof. Let M and N be the negative mixed crossings. Let $\tilde{a}$ be the arc on $\tilde{\ell}_1$ from M to N and $\tilde{b}$ the arc on $\tilde{\ell}_1$ from N to M . Let $\tilde{c}$ be the arc on $\tilde{\ell}_2$ from M to N and $\tilde{d}$ the arc on $\tilde{\ell}_2$ from N to M . Suppose that there is a self-crossing of $\tilde{\ell}_1$ between $\tilde{a}$ and $\tilde{b}$. Then by Lemma 2.2 we have the result. Similarly if there is a self-crossing of $\tilde{\ell}_2$ between $\tilde{c}$ and $\tilde{d}$ then we have the result. Therefore we may suppose that there are no such self-crossings. We may suppose without loss of generality that $\tilde{a}$ and $\tilde{c}$ has a positive mixed crossing. Using Lemma 2.2 we may suppose that $\tilde{d}$ is a simple arc. First suppose that $\tilde{a}$ has a positive mixed crossing on $\tilde{d}$. Then we have the trivial link by changing the crossings without changing M and N as illustrated in Fig 4.69. Next suppose that $\tilde{a}$ has no mixed crossings on $\tilde{d}$ except M and N .

Then we change the crossings of $\tilde{L}$ so that $\tilde{b}$ is parallel to $\tilde{d}$ and apply Lemma 2.7 on the tangle diagram that is complementary to a neighborhood of $\tilde{d}$ and we have the trivial link. See Fig. 4.70. $\square$




FIGURE 4.69.




FIGURE 4.70.

Lemma 4.10. *Let $\tilde{L} = \tilde{\ell}_1 \cup \tilde{\ell}_2 \cup \tilde{\ell}_3$ be a 2-almost positive diagram of a 3-component link L . Suppose that $\tilde{\ell}_1 \cap \tilde{\ell}_2$ has the two negative mixed crossings and has no other mixed crossing. Suppose that $\tilde{\ell}_3$ has mixed crossings with both of $\tilde{\ell}_1$ and $\tilde{\ell}_2$. Then L is greater than or equal to the link of Fig. 1.4 (d).*

Proof. If we find a part as illustrated in Fig. 4.71 (a) in $\tilde{L}$ then we have the link of Fig. 1.4 (d). If there are no such parts then we have one of the parts illustrated in Fig. 4.71 (b). Then we have the link of Fig. 1.4 (d) or the link L_0 in Fig. 4.71 (b) that is greater than or equal to the link of Fig. 1.4 (d) as illustrated in Fig. 4.71 (c). $\square$

Theorem 4.11. *Let $\tilde{L}$ be a 2-almost positive diagram of a link L . Then*

- (1) $L \geq$ right-handed trefoil knot (plus trivial components) or




FIGURE 4.71.

- (2) $L \geq 6_2$ (Fig. 1.4 (a)) (plus trivial components) or
- (3) $L \geq$ right-handed Hopf link (plus trivial components) or
- (4) $L \geq$ disjoint or connected sum of right-handed trefoil knot and left handed Hopf link (plus trivial components) or
- (5) $L \geq$ Whitehead link (Fig. 1.4 (b)) (plus trivial components) or
- (6) $L \geq$ disjoint and/or connected sum of two right-handed Hopf links and a left-handed Hopf link (plus trivial components) or
- (7) $L \geq$ disjoint or connected sum of $(2,4)$ -torus link (Fig. 1.4 (c)) and a left-handed Hopf link (plus trivial components) or
- (8) $L \geq$ the link of Fig. 1.4 (d) (plus trivial components) or
- (9) $\tilde{L}$ is a diagram obtained from the diagrams in Fig. 1.2 and Fig. 1.6 and their reversals on $\mathbb{S}^2 = \mathbb{R}^2 \cup \infty$ (reflection in y axis) by performing, possibly, diagram-disjoint sum operation, diagram-connected sum operation, and first and second Reidemeister moves which increase the number of crossings, or
- (10) $\tilde{L}$ is a diagram obtained from $\tilde{L}_1$, $\tilde{L}_2$ or $\tilde{L}_3$ of Fig. 4.72 by performing, possibly, first Reidemeister moves which increase the number of crossings and adding trivial circles, or
- (11) $\tilde{L}$ is a diagram obtained from $\tilde{L}_3$ of Fig. 4.72 and $\tilde{T}_1$ or $\tilde{T}_2$ of Fig. 4.73 by, possibly, performing first Reidemeister moves which increase the number of crossings, diagram-disjoint sum operation or diagram-connected sum operation (plus almost trivial components).

Proof. We decompose $\tilde{L}$ into prime factors. If the two negative crossings belong to different prime factors then the result follows by Theorem 3.2. Therefore it is sufficient to consider the case that $\tilde{L}$ is prime and has just two negative crossings, say M and N . Then the case that $\tilde{L}$ is not prime immediately follows.

Case 1. M and N are self-crossings of a component, say $\tilde{\ell}_1$ of $\tilde{L}$.




FIGURE 4.72.




FIGURE 4.73.

We first note that by applying Lemma 2.2 we have that $\tilde{\ell}_1$ is greater than or equal to a trivial knot. Therefore if $\tilde{L}$ has other components then we have $L \geq$ right-handed Hopf link. Hence we may suppose $\tilde{L} = \tilde{\ell}_1$. We have the following three cases according to the position of M and N as illustrated in Fig. 4.74




FIGURE 4.74.

First suppose $(\tilde{a} \cup \tilde{b}) \cap (\tilde{c} \cup \tilde{d}) \neq \emptyset$. Let P be a crossing of $(\tilde{a} \cup \tilde{b}) \cap (\tilde{c} \cup \tilde{d})$. Let $\tilde{T}$ be the complementary tangle diagram of $\tilde{L}$ at P . Then $\tilde{T}$ is a prime R2-reduced 2-string tangle diagram with vertical connection with just two negative crossings, one is a mixed crossing and the other is a self-crossing. Note that the X_+ -closures of the tangles $T(-2), T(3_1, 0_1), T(0_1, 3_1)$ and T_2 in Fig. 4.7 are right-handed trefoil knots and the X_+ -closures of the tangles T_1, T_3 and T_4 in Fig. 4.7 are 6_2 . The X_+ -closures of the tangles T_5 and

T_6 in Fig. 4.7 are greater than or equal to the X_- -closures of the tangles T_5 and T_6 in Fig. 4.7 respectively and these knots are 6_2 . Note also that the X_+ -closures of the tangle diagrams $\tilde{T}_1, \tilde{T}_{2++}, \tilde{T}_{2+-}, \tilde{T}_{2-+}, \tilde{T}_{2--}, \tilde{T}_{3+}$ and $\tilde{T}_{3-}$ in Fig. 4.8 are not R2-reduced. The X_+ -closures of the tangle diagrams $\tilde{T}_{4+}$ and $\tilde{T}_{5-}$ in Fig. 4.8 are equal to the link diagram $\tilde{T}_2$ of Fig. 1.6. The X_+ -closures of the tangle diagrams $\tilde{T}_{4-}$ and $\tilde{T}_{5+}$ in Fig. 4.8 are equal to the reversal of the link diagram $\tilde{T}_2$ of Fig. 1.6. The X_+ -closures of the tangle diagrams $\tilde{T}_{6+}$ and $\tilde{T}_{7-}$ in Fig. 4.8 are equal to the link diagram $\tilde{T}_1$ of Fig. 1.6. The X_+ -closures of the tangle diagrams $\tilde{T}_{6-}$ and $\tilde{T}_{7+}$ in Fig. 4.8 are equal to the reversal of the link diagram $\tilde{T}_1$ of Fig. 1.6. Then by Lemma 4.2 we have $L \geq$ right-handed trefoil knot or $L \geq 6_2$ if and only if $\tilde{L}$ is not equal to the link diagrams $\tilde{T}_1$ or $\tilde{T}_2$ of Fig. 1.6 or their reversals (reflections in y axis).

Next suppose $(\tilde{a} \cup \tilde{b}) \cap (\tilde{c} \cup \tilde{d}) = \emptyset$. Choose disjoint disks D_1 and D_2 on $\mathbb{S}^2$ such that D_1 contains $\tilde{a}$ and $\tilde{b}$ and does not contain M and N , D_2 contains $\tilde{c}$ and $\tilde{d}$ and does not contain M and N . See Fig. 4.75. In Case 1.1 we apply Lemma 4.4 to D_1 and Lemma 2.9 to D_2 . Note that if we have the tangle $T(3_1, 0_1)$ or $T(0_1, 3_1)$ on D_1 then by trivializing the tangle on D_2 we have the right-handed trefoil knot. If we have the tangle $R(T(-3, -1))$ on D_1 then by trivializing the tangle on D_2 we have 6_2 . If we have the tangle $T(-1, -2)$ on D_2 then by trivializing the tangle on D_1 we have the right-handed trefoil knot. Then taking the fact that $\tilde{T}$ is R2 reduced into account we have that $L \geq$ right-handed trefoil knot or $L \geq 6_2$ if and only if $\tilde{L}$ is not $\tilde{T}_1$ in Fig. 4.72. In Case 1.2 we have the right-handed trefoil knot by applying Lemma 2.10 to D_2 and trivializing the tangle on D_1 .

In Case 1.3 we apply Lemma 4.6 to D_1 and Lemma 4.1 to D_2 . If we have one of the tangles in Fig. 4.56 except $T(1/2, -3)$ on D_1 then by trivializing the tangle on D_2 we have the right-handed trefoil knot. If we have the tangle $T(1/2, -3)$ on D_1 then by trivializing the tangle on D_2 we have 6_2 . If we have one of the tangles in Fig. 4.1 on D_2 then by taking the positive tangle with one crossing on D_1 we have the right-handed trefoil knot. Therefore, because $\tilde{L}$ is R2-reduced, the diagram on D_1 is a positive diagram whose underlying projection is one of the projections $\hat{T}_1, \hat{T}_3, \hat{T}_4$ and $\hat{T}_5$ in Fig. 4.57 or their flypes. Also because $\tilde{L}$ is R2-reduced, we have that the diagram on D_2 is a positive diagram whose underlying projection is $\hat{T}(1, 2n)$ or $\hat{T}(2n, 1)$ in Fig. 4.2 unless it is a trivial diagram. Suppose that it is not a trivial diagram. Then by Lemma 2.7 and Lemma 2.10 we have the right-handed trefoil knot unless the underlying projection of the diagram on D_1 is $\hat{T}_1$ in Fig. 4.57. See Fig. 4.76. If the underlying projection of the diagram on D_1 is $\hat{T}_1$ in Fig. 4.57 then we either have the right-handed trefoil knot or the diagram $\tilde{T}_9$ in Fig. 1.6. See Fig. 4.77. Suppose that the diagram on D_2 is trivial. Then we have that $\tilde{L}$ is (a reversal of) $\tilde{T}_3, \tilde{T}_4, \tilde{T}_5, \tilde{T}_6, \tilde{T}_7, \tilde{T}_8$ or $\tilde{T}_{10}$ in Fig. 1.6 or $\tilde{L}_2$ in Fig. 4.72.

Case 2. M and N are self-crossings of different components of $\tilde{L}$.

In this case we easily have $L \geq$ right-handed Hopf link (plus trivial components).




FIGURE 4.75.




FIGURE 4.76.




FIGURE 4.77.

Case 3. One of M and N , say M , is a self-crossing of a component, say $\tilde{\ell}_1$, and N is a mixed crossing of components, say $\tilde{\ell}_i$ and $\tilde{\ell}_j$.

Then we have either $L \geq$ right-handed Hopf link (plus trivial components) or $\tilde{L} = \tilde{\ell}_i \cup \tilde{\ell}_j$. If $\tilde{L} = \tilde{\ell}_i \cup \tilde{\ell}_j$ then choose a small disk as illustrated in Fig. 4.78 and apply Lemma 4.2 to the tangle diagram on the complementary disk on $\mathbb{S}^2$. Note that because $\tilde{L}$ is R2-reduced, we have no tangle diagrams in Fig. 4.8. Then we have $L \geq$ right-handed trefoil knot and an unknot or $L \geq$ right-handed Hopf link or $L \geq$ Whitehead link.

Case 4. Both M and N are mixed crossings.

First suppose that $\tilde{L}$ has four or more components. Then using Lemma 4.9 we have (3) or (6). Suppose that $\tilde{L}$ has three components, say $\tilde{\ell}_1, \tilde{\ell}_2$




FIGURE 4.78.

and $\tilde{\ell}_3$. Suppose that M and N are mixed crossings of different pair of components. We may suppose without loss of generality that M is a mixed crossing between $\tilde{\ell}_1$ and $\tilde{\ell}_2$ and N is between $\tilde{\ell}_2$ and $\tilde{\ell}_3$. Suppose that $\tilde{\ell}_1$ and $\tilde{\ell}_2$ has four or more mixed crossings between them. Then by applying Lemma 2.10 to a complementary tangle diagram of $\tilde{\ell}_1 \cup \tilde{\ell}_2$ at M we have that the diagram $\tilde{\ell}_1 \cup \tilde{\ell}_2$ is greater than or equal to the right-handed Hopf link. Then we have (3). Therefore we have that $\tilde{\ell}_1$ and $\tilde{\ell}_2$ have just two mixed crossings between them. Then by the primeness of $\tilde{L}$ we have that $\tilde{\ell}_1$ and $\tilde{\ell}_3$ has mixed crossings between them. Note that there is a mixed crossing, say P between $\tilde{\ell}_1$ and $\tilde{\ell}_3$ such that $\tilde{\ell}_1$ is over $\tilde{\ell}_3$ at P , and there is another mixed crossing, say Q between $\tilde{\ell}_1$ and $\tilde{\ell}_3$ such that $\tilde{\ell}_1$ is under $\tilde{\ell}_3$ at Q . If $\tilde{\ell}_2$ is both over $\tilde{\ell}_1$ at M and over $\tilde{\ell}_3$ at N , or both under $\tilde{\ell}_1$ at M and under $\tilde{\ell}_3$ at N , then we have (3). Suppose that $\tilde{\ell}_2$ is over $\tilde{\ell}_1$ at M and under $\tilde{\ell}_3$ at N . Then we change crossings of $\tilde{L}$ so that $\tilde{\ell}_1$ is under everything except at P , and $\tilde{\ell}_3$ is over everything except at P . Then without changing M and N we have a right-handed Hopf link. The other case is similar. Next suppose that M and N are between the same pair of components, say $\tilde{\ell}_1$ and $\tilde{\ell}_2$. If $\tilde{\ell}_1$ and $\tilde{\ell}_2$ have four or more mixed crossings between them then we have (3) by Lemma 4.9. Thus we have that $\tilde{\ell}_1$ and $\tilde{\ell}_2$ have no other mixed crossings between them. Then by the primeness of $\tilde{L}$ and by Lemma 4.10 we have (8). Suppose that $\tilde{L}$ has just two components. Then using Lemma 4.8 to the complementary tangle diagram of M we have (3), (4) or (5) if $\tilde{L}$ is not $\tilde{T}_{11}$ of Fig. 1.6 nor $\tilde{L}_3$ of Fig. 4.72. This completes the proof. $\square$

Theorem 1.8, Theorem 1.9 and Corollary 1.10 are immediate corollaries of Theorem 4.11.

5. 3-ALMOST POSITIVE KNOTS

Theorem 5.1. *Let K be a 3-almost positive knot. Then either $K \geq$ trivial knot or K is the left-handed trefoil knot (plus positive knots as connected summands).*

Proof. Let $\tilde{K}$ be a diagram of K with three negative crossings N_1 , N_2 and N_3 . Suppose for example that a loop from N_1 to N_1 passes through at most one of N_2 and N_3 , or passes through both of them as over-crossing, or passes through both of them as under-crossing. Then by Lemma 2.2 we have that K is greater than or equal to the trivial knot. Thus we are left with the case in which every loop from N_i to N_i passes through two other negative crossings, one as over-crossing the other as under-crossing. This situation is illustrated in Fig. 5.1.





FIGURE 5.1.

We name the points as in Fig. 5.1. Let $\tilde{a}_i = A_i B_i$. We consider the suffix modulo six. If $\tilde{a}_i \cap \tilde{a}_{i+1}$ is nonempty for some i , then it reduces to the case with two negative crossings as any crossing of $\tilde{a}_i \cap \tilde{a}_{i+1}$ can serve as a crossing P from Lemma 2.2. Suppose $\tilde{a}'_i \cap (\tilde{a}_{i+2} \cup \tilde{a}_{i-2})$ is nonempty (the spine $\tilde{a}'_i$ of $\tilde{a}_i$ has been defined in Section 2.1). We may assume without loss of generality that $\tilde{a}'_i \cap \tilde{a}_{i+2}$ is nonempty. Let P be the first crossing starting from A_{i+2} with $\tilde{a}_i$. We reduce $A_{i+2}P$ and $\tilde{a}_{i+1}$ to become simple arcs using Lemma 2.2. Then we pull down $PB_{i+2}A_{i+3}B_{i+3}$, pull up $\tilde{a}_{i+4}$ and we have a knot with no negative crossings as illustrated in Fig. 5.2 and the case is completed.

FIGURE 5.2. Reduction in the case of $\tilde{a}'_i \cap \tilde{a}_{i+2} \neq \emptyset$

Now suppose $\tilde{a}'_i \cap \tilde{a}_{i+3}$ is nonempty. We simplify $\tilde{a}_{i+1}$ and $\tilde{a}_{i+2}$. Let P be a crossing of $\tilde{a}'_i \cup \tilde{a}_{i+3}$. We pull down $A_{i+3}P$ and pull up $PB_{i+3}A_{i+4}B_{i+4}$. Then we have a diagram with only one negative crossing as illustrated in Fig. 5.3.

Therefore we may suppose that no $\tilde{a}'_i$ has crossings with other $\tilde{a}_j$'s. Then we have that the union of all $\tilde{a}'_i$ and $B_i A_{i+1}$ forms a standard diagram of a left-handed trefoil knot. Now suppose $\tilde{a}_i \cap \tilde{a}_j \neq \emptyset$. Then the result follows by Lemma 2.7 as illustrated in Fig. 5.4.




FIGURE 5.3.




FIGURE 5.4.

Therefore $\tilde{a}_i \cap \tilde{a}_j$ is empty for all $i \neq j$. Then the knot K is the left-handed trefoil knot (possibly with some positive knots connected summed). This completes the proof. $\square$

Corollary 1.12 follows immediately from Theorem 1.11 and the fact that the connected sum of the left-handed trefoil knot and a nontrivial positive knot has non-positive signature.

We do not know whether any knot with the left-handed trefoil knot as a connected sum summand can dominate the trivial knot. We think it is highly unlikely.

6. COROLLARIES

In the previous sections we have been applying our “domination” technique mainly to the classical (Trotter-Murasugi) signature, $\sigma(L)$. Here we give several other applications of the theory including amphicheirality, sliceness, bounds on Jones polynomial, and Tristram-Levine signatures.

- Corollary 6.1.** (a) *A nontrivial almost positive link is not amphicheiral.*
 (b) *2-almost positive link L without a trivial component is amphicheiral if and only if either*
 (i) *L is a connected or disjoint sum of right-handed and left-handed Hopf links, or*
 (ii) *L is the figure eight knot.*

Proof. If L is amphicheiral then the signature $\sigma(L) = 0$, so (a) follows from Corollary 1.7. To prove (b), we apply Corollary 1.10. The cases (1) and (3) of Corollary 1.10 have $\sigma(L) = 0$. An amphicheiral link L has the global linking number $lk(L) = 0$. Among the links described in Corollary 1.10 only connected or disjoint sums of right and left-handed Hopf links (plus possibly trivial components) have $lk(L) = 0$, and they are amphicheiral. For a twist knot it is well known by the classification of 2-bridge knots [Sch] that it is amphicheiral if and only if it is the trivial knot or the figure eight knot. $\square$

- Corollary 6.2.** (a) *A nontrivial almost positive link is not a slice link.*
 (b) *A nontrivial 2-almost positive link is a (smoothly) slice link if and only if it is the Stevedore knot (possibly with additional trivial components).*

Proof. If L is a slice link then $\sigma(L) = 0$ and any pair of components have linking number zero. Therefore (a) follows from Corollary 1.7 and (b) from Corollary 1.10 by observing that a twist knot is smoothly slice if and only if it is the trivial knot or the stevedore knot (Casson and Gordon [Ca-G]). $\square$

Recall that the Jones polynomial [Jo] of a link L is an element of the ring of the Laurent polynomials in variable $\sqrt{t}$. That is $V_L(t) \in \mathbb{Z}[\sqrt{t}^{\pm 1}]$ satisfying the skein relation:

$$t^{-1}V_{L_+}(t) - tV_{L_-}(t) = (\sqrt{t} - \frac{1}{\sqrt{t}})V_{L_0}(t)$$

and normalized to be 1 at the trivial knot.

Let $d_{\min}V_L(t)$ denote the lowest power of t in $V_L(t)$. Notice that $V_L(t) \in \mathbb{Z}[t^{\pm 1}]$ if the number of components of L , $\mu(L)$, is odd and $\sqrt{t}V_L(t) \in \mathbb{Z}[t^{\pm 1}]$ if $\mu(L)$ is even, so the inequalities in Corollary 6.3 (a)-(c) are sharp in the case of a knot.

Corollary 6.3. *Let L be a nontrivial nonsplit link, then*

- (a) *If L is positive then $d_{\min}V_L(t) \geq 3/2$ unless*
 (i) *L is a $(2, 2k)$ -torus link with anti-parallel orientation of components (Fig. 1.5) (denote it $T_{2,2k}^{an}$) then $d_{\min}V_L(t) = 1/2$ or,*
 (ii) *$L = T_{2,2k}^{an} \# T_{2,2k}^{an}$; then $d_{\min}V_L(t) = 1$ or,*
 (iii) *L is a pretzel knot $L(p_1, p_2, p_3)$ of Fig. 1.1 or a 3-component pretzel link $L(q_1, q_2, q_3)$ of Fig. 2.55 (c); then $d_{\min}V_L(t) = 1$.*
 (b) *If L is an almost positive link then $d_{\min}V_L(t) \geq 1/2$.*
 (c) *If L is a 2-almost positive link then $d_{\min}V_L(t) \geq -3/2$ unless either*
 (i) *L is a left-handed Hopf link (denoted H_-) then $d_{\min}V_L(t) = -5/2$ or,*

- (ii) $L = H_- \# T_{2,2k}^{an}$ then $d_{\min} V_L(t) = -2$ or,
- (iii) L is a twist knot for which $d_{\min} V_L(t) = -2$.
- (d) If K is a 3-almost positive knot then $d_{\min} V_L(t) \geq -3$ except the left-handed trefoil knot for which $d_{\min} V_L(t) = -4$.

Proof. K. Murasugi showed ([M-3] Theorem 13.3) that if D is a diagram of a non-split link L with $c_-(D)$ negative crossings then

$$d_{\min} V_L(t) \geq -c_-(D) - \frac{1}{2}\sigma(L).$$

Murasugi's inequality and Corollary 1.12 (a) give Corollary 6.3 (d). Murasugi's inequality and Corollary 1.10 give Corollary 6.3 (c) (there is no need to perform calculations for exceptional cases (i)-(iii) because for alternating diagrams (or their connected sums) Murasugi's inequality becomes the equality ([M-3])). Murasugi's inequality and Corollary 1.3 suffice to prove (a). In the case (b) Murasugi's inequality and the fact that $\sigma(L) < 0$ are not sufficient (we get only $d_{\min} V_L(t) \geq -1/2$; or in the case of knots $d_{\min} V_L(t) \geq 0$) and to improve this we would have to show that Murasugi's inequality is a strict inequality in our case (an almost positive diagram with alternating connected summands has a nugatory crossing). We choose, however, a different method which is of interest on its own and generalizes Corollary 6.3 (a) and (b).

Theorem 6.4. *Let D be a diagram of an oriented link L , then*

- (a) *If D is positive then*

$$d_{\min} V_L(t) = \frac{1}{2}(c(D) - s(D) + 1)$$

where $c(D)$ is the number of crossings of D and $s(D)$ is the number of Seifert circles of D . In particular, if L is a knot then $d_{\min} V_L(t) = g(D)$, where $g(D)$ is the genus of the Seifert surface obtained from D by the Seifert algorithm⁷.

- (b) *If D has one negative crossing say P then either*
 - (i) *P is a singular crossing (i.e. there are no other crossings joining the same, as P , Seifert circles of D) then*

$$d_{\min} V_L(t) = \frac{1}{2}(c(D) - s(D) + 1)$$

or

- (ii) *P is not a singular crossing and then*

$$d_{\min} V_L(t) = \frac{1}{2}(c(D) - s(D) + 1) - 1 = \frac{1}{2}(c_+(D) - c_-(D) - s(D) + 1)$$

In particular Corollary 6.3 (b) follows.

□

To prove Theorem 6.4 we need the following technical Lemma.

Lemma 6.5. *Let D be an oriented link diagram and $F = P_1, P_2, \dots, P_k$ a family of different crossings of D .*

⁷After the first version of the paper was written, Kronheimer and Mrowka [KM] and then Rasmussen [Ras] proved that $g(D)$ is equal to genus of the knot L as well as a slice genus of L .

- (a) Assume that A -splittings $\left(\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \rightarrow \begin{array}{c} \diagup \diagup \\ \diagdown \diagdown \end{array} \right)$ of crossings $P_1, P_2, \dots, P_k$ produce a diagram D' such that two arcs obtained from the splitting of P_i , for any i , are on different connected components of D' considered as a graph (we say that D is $+$ -adequate with respect to crossings F). Then

$$4d_{\min} V_D(t) = 4d_{\min} V_{D'}(t) + 3(c_+(D) - c_-(D) - (c_+(D') - c_-(D')) - c(F))$$

where $c(F)$ is the number of crossings in the family F .

- (b) Assume that all crossings in F are positive and D' is obtained from D by smoothing $\left(\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \rightarrow \begin{array}{c} \diagup \diagup \\ \diagdown \diagdown \end{array} \right)$ all crossings of F , and for any i two arcs obtained from the smoothing of P_i are on different connected components of D' considered as a graph. Then

$$d_{\min} V_D(t) = d_{\min} V_{D'}(t) + \frac{1}{2}c(F)$$

- (c) (compare [M-P-1]) Let $D = D_1 * D_2$ (planar star (Murasugi) product) and D_2 be a positive diagram. Then

$$d_{\min} V_D(t) = d_{\min} V_{D_1}(t) + d_{\min} V_{D_2}(t) = d_{\min} V_{D_1}(t) + \frac{1}{2}(c(D_2) - s(D_2) + 1).$$

Proof. We use the Kauffman bracket approach to $V_L(t)$.

Recall that the Kauffman bracket polynomial of a link diagram, $\langle D \rangle \in Z[A^{\pm 1}]$, satisfies the Kauffman bracket skein relations [Ka-2]:

$$\langle D_+ \rangle = A \langle D_0 \rangle + A^{-1} \langle D_\infty \rangle, \quad \langle D \sqcup O \rangle = (-A^2 - A^{-2}) \langle D \rangle,$$

and is normalized to be 1 at the crossingless diagram of the unknot. The Jones polynomial $V_L(t) \in Z[t^{\pm \frac{1}{2}}]$ can be obtained from the Kauffman bracket polynomial of any oriented diagram D of a link L by putting $t = A^{-4}$ in $V_L(t) = (-A^3)^{-w(D)} \langle D \rangle$, where $w(D)$ is the writhe or Tait number of an oriented diagram D (that is, $w(D) = c_+(D) - c_-(D) = \sum_p \text{sgn}(p)$ where the sum is taken over all crossings p of oriented diagram D).

- (a) We can repeat essentially the proofs of [K], [L-T] and [T] (see for example the proof of Proposition 1 in [L-T] about the maximal and minimal terms of the bracket $\langle D \rangle$ of an adequate diagram D). In particular, we get immediately that A -smoothing along F contributes to the maximal term in $\langle D \rangle$, so $d_{\max} A \langle D \rangle = d_{\max} A \langle D' \rangle + c(F)$, and thus $-4d_{\min} V_D(t) + 3w(D) = -4d_{\min} V_{D'}(t) + 3w(D') + c(F)$.
- (b) Observe that a smoothing at a positive crossing is the same as A -splitting the crossing so (b) follows from (a) (Figure 6.1 illustrates this fact).




FIGURE 6.1. A -splitting agrees with smoothing for a positive crossing

- (c) The family of crossings of D_2 satisfies the assumptions of (b) so (c) follows from (b).

□

PROOF OF THEOREM 6.4:

- (a) We apply Lemma 6.5 (a) to family F of all crossings of D , then D' is a collection of all Seifert circles of D so $d_{\min} V_{D'}(t) = 1/2(-s(D) + 1)$. Therefore $d_{\min} V_D(t) = 1/2(c(D) - s(D) + 1)$.
- (b) (i) P is a singular crossing and F a family of all other crossings of D . Let D_+^P be a positive diagram obtained from D by changing the crossing P from negative to positive. Finally let D' (resp. $(D_+^P)'$) denote the diagram obtained from D (resp. D_+^P) by smoothing all crossings from family F . Notice that D' represents the link ambient isotopic to $(D_+^P)'$ and in both cases we can use Lemma 6.5 (b) and family F (see Fig. 6.2).




FIGURE 6.2. P is a nugatory crossing in D' and $(D_+^P)'$

Therefore

$$d_{\min} V_D(t) = d_{\min} V_{D'}(t) + \frac{1}{2}c(F) = d_{\min} V_{(D_+^P)'}(t) + \frac{1}{2}c(F) = d_{\min} V_{D_+^P}(t)$$

Furthermore D_+^P is a positive diagram so by part (a) of Theorem 6.4

$$d_{\min} V_{D_+^P}(t) = \frac{1}{2}(c(D_+^P) - s(D_+^P) - 1).$$

The formula from (b)(i) holds because $c(D_+^P) = c(D)$ and $s(D_+^P) = s(D)$.

- (ii) Let $Q_1, Q_2, \dots, Q_n$ be all other crossings of D which join the same Seifert circles at P . Let F be the set of crossings of D different from P and $Q_1, Q_2, \dots, Q_n$. Furthermore let $D_0 = D_{00}^{PQ_1}$ denote the diagram obtained from D by smoothing crossings P and Q_1 . Finally let D' (resp. D'_0) denote the diagram obtained from D (resp. D_0) by smoothing all crossings from family F . Notice that D' represents a link isotopic to D'_0 and in both cases we can apply Lemma 6.5 (b) using family F (see Fig. 6.3).

Therefore (as in Case (i)), $d_{\min} V_D(t) = d_{\min} V_{D_0}(t)$. Also D_0 is a positive diagram so by part (a) of Theorem 5.5,

$$d_{\min} V_{D_0}(t) = \frac{1}{2}(c(D_0) - s(D_0) - 1).$$




FIGURE 6.3. P is a negative crossing which is non-singular and cancels in D'

Because $c(D) = c(D_0) + 2$, $s(D) = s(D_0)$, we get the formula from Theorem 6.4 (b)(ii).

Now assume that D is a connected diagram, then $c(D) - s(D) + 1 \geq 0$, and the equality holds if and only if all crossings are nugatory so if L is nontrivial then $c(D) - s(D) + 1 \geq 1$. Now assume additionally that P is the only negative crossing of D , P is not singular and D has no nugatory crossings. If

- (1) $c(D) - s(D) + 1 = 1$ then D looks as in Fig. 6.4 (i) so it represents a split link.
- (2) $c(D) - s(D) + 1 = 2$ then D looks as in Fig. 6.4 (ii) or (iii) so it represents the trivial knot or a split link.

Corollary 6.3 (b) follows.




FIGURE 6.4. $c(D) - s(D) + 1 = 1$ or $c(D) - s(D) + 1 = 2$ with a non-singular negative crossing

Remark 6.6. *If we consider the Seifert graph associated to D , say $\Gamma(D)$ (see [Cr-Mo] or [M-P-2]) then $c(D) - s(D) + 1$ is the first Betti number of $\Gamma(D)$ (called cyclomatic number in graph theory). Our method is to analyze diagrams such that $\Gamma(D)$ has small cyclomatic number. In essence we can then extend Corollary 6.3 (c) to $d_{\min} V_L(t) \geq 1/2$ unless... But the list of exceptions would be rather long so not very interesting (unless we will try to use this to determine which links are 2-almost positive).*




FIGURE 6.5. The Seifert graph in (i) is the graph of Fig. 6.4 (ii) for n odd. The Seifert graph in (ii) is the graph of Fig. 6.4 (ii) for n even or of Fig. 6.4 (iii)

Corollary 6.7. *A nontrivial positive link L has unknotting number one if and only if L is a right-handed Hopf link (plus, possibly, trivial components) or L is a (positive) twist knot (possibly with additional trivial components).*

Proof. A Hopf link and twist knots have unknotting number one, we will show the converse. Let $u(L)$ be the unknotting number of L . Then $2u(L) \geq |\sigma(L)|$ [M-1] and clearly $u(L) \geq \sum_{i < j} |lk(\ell_i, \ell_j)|$ where the sum is taken over all different pairs of components of L . Therefore by Corollary 1.9 we are left with pretzel knots $L(p_1, p_2, p_3)$ of Fig. 1.1. These pretzel knots bound genus one Seifert surfaces in a natural manner. Since genus one unknotting number one knots are doubled knots [Ko], [S-T] and pretzel knots are simple [Ko], we are left with twist knots. $\square$

In the next Theorem we use our domination theorems to show, generalizing [P-1], that Tristram-Levine signature satisfies some inequalities for k -almost positive links.

Let $\sigma_\psi(L)$ be the Tristram-Levine signature of L ; we assume $|\psi| = 1$. We use the convention that $\sigma_\psi(L)$ is the signature of the Hermitian matrix $H = \bar{\psi}A_L + \psi A_L^T$, where A_L is a Seifert matrix of L . Because for $Re(\psi) \geq 0$ we have $\sigma_\psi(L_+) \leq \sigma_\psi(L_-)$ ([P-Tr-1, P-2, P-3]) we can use domination results: Theorems 2.15, 3.2, 4.11 and 5.1 to bound Tristram-Levine signature of positive, almost positive, 2-almost positive and 3-almost positive links. In particular, we have.

Theorem 6.8. (i) ([P-1]): *If L is a nontrivial positive link and $Re(\psi) > \frac{1}{2}$, then $\sigma_\psi(L) < 0$.*

(ii) *If K is a nontrivial positive knot and $Re(\psi) \geq 0$ then $\sigma_\psi(K) \leq \sigma_\psi(\bar{5}_1)$ or K is a connected sum of some pretzel knots of type $L(2k_1 + 1, 2k_2 + 1, 2k_3 + 1)$, where $k_i \geq 0$; see Figure 1.1. Furthermore we have (compare e.g. Examples 5.16, and 5.14 of [P-3]):*

$$\sigma_\psi(\bar{5}_1) = \begin{cases} -4 & \text{if } Re(\psi) > \frac{1+\sqrt{5}}{4} \\ -3 & \text{if } Re(\psi) = Re(e^{\pi i/5}) = \frac{1+\sqrt{5}}{4} \approx 0.809... \\ -2 & \text{if } Re(e^{2\pi i/5}) \leq Re(\psi) < Re(e^{\pi i/5}) \\ -1 & \text{if } Re(\psi) = Re(e^{2\pi i/5}) = \frac{\sqrt{5}-1}{4} \approx 0.309... \\ 0 & \text{if } 0 \leq Re(\psi) < Re(e^{2\pi i/5}). \end{cases}$$

For $1 + k_1 + k_2 + k_3 + k_1k_2 + k_1k_3 + k_2k_3 > 0$,

$$\sigma_\psi(L(2k_1+1, 2k_2+1, 2k_3+1)) = \begin{cases} -2 & \text{if } Re(\psi) > \frac{1}{2\sqrt{1+k_1+k_2+k_3+k_1k_2+k_1k_3+k_2k_3}} \\ -1 & \text{if } Re(\psi) = \frac{1}{2\sqrt{1+k_1+k_2+k_3+k_1k_2+k_1k_3+k_2k_3}} \\ 0 & \text{if } 0 \leq Re(\psi) < \frac{1}{2\sqrt{1+k_1+k_2+k_3+k_1k_2+k_1k_3+k_2k_3}}. \end{cases}$$

In the case that $k_1 = k_2 = k_3 = 0$ represents the right handed trefoil knot, $\bar{3}_1$.

(iii) If L is a nontrivial almost positive link and $Re(\psi) \geq 0$ then

$\sigma_\psi(L) \leq \sigma_\psi(\bar{3}_1)$ or $\sigma_\psi(L) \leq \sigma_\psi(H_+)$, and if L has a nontrivial component and $Re(\psi) > \frac{1}{2}$, we have $\sigma_\psi(L) \leq -2$

(iv) If K is a nontrivial 2-almost positive knot and $Re(\psi) \geq 0$ then,

$\sigma_\psi(K) \leq \sigma_\psi(\bar{3}_1)$ or $\sigma_\psi(K) \leq \sigma_\psi(6_2)$ or K is a twist knot with a negative clasp (see Figure 1.3). We have the Tristram-Levin signature equal to zero in the last case, furthermore for $\sigma_\psi(6_2)$ we have⁸:

$$\sigma_\psi(6_2) = \begin{cases} -2 & \text{if } Re(\psi) > \frac{1}{2}\sqrt{\frac{1+\sqrt{5}}{2}} \\ -1 & \text{if } Re(\psi) = \frac{1}{2}\sqrt{\frac{1+\sqrt{5}}{2}} \approx 0.636... \\ 0 & \text{if } 0 \leq Re(\psi) < \frac{1}{2}\sqrt{\frac{1+\sqrt{5}}{2}}. \end{cases}$$

(v) If K is a 3-almost positive knot, and $Re(\psi) \geq 0$ then

$\sigma_\psi(K) \leq 0$ or K is the left handed trefoil knot.

In [P-2] we observe that domination results can be applied to any signature-like invariant σ satisfying Trotter type inequalities. In 1987 or 1991 we could only speculate about existence of such invariants⁹ [P-1, P-Tr-3].

Finally, we prove Theorem 1.13(b) following Cochran and Gompf paper [Co-G], using Theorem 1.8, and the computation by S. Akbulut (compare Problem 4.2 of [Kir]).

Theorem 1.13 (b). If K is a 2-almost positive knot different from a twist knot with a negative clasp then $K(1/n)$ (i.e. $1/n$ surgery on K , $n > 0$) is a homology 3-sphere that does not bound a compact, smooth homology 4-ball. Furthermore, $K(1/n)$ has a nontrivial Floer homology.

Proof. It is proved in [Co-Li, Co-G] that if $K_1 \geq K_2$ then $K_1 \succeq K_2$. Furthermore, Cochran and Gompf prove in [Co-G] that if $K_1 \succeq K_2$ and $K_2(1/1)$ is a homology sphere which bounds NSPD (non-standard positive definite) homology sphere, then for any $n > 0$, the homology sphere $K_1(1/n)$ bounds NSPD homology sphere. Then, using Donaldson result they conclude that $K_1(1/n)$ does not bound a compact, smooth homology 4-ball¹⁰. The fact that $\bar{3}_1(1/1)$ bounds NSPD homology sphere is an important tool in [Co-G]. Because our generalization of Cochran-Gompf result uses Theorem 1.8 we need to show that $6_2(1/1)$ bounds NSPD homology sphere. The proof of this fact was provided to us by S. Akbulut (in letter sent February

⁸In the convention of [Gor, Ch-L] one defines the Tristram-Levine signature function of variable ξ ($|\xi| = 1$) as $\sigma_L(\xi) = \sigma((1 - \bar{\xi})A + (1 - \xi)A^T)$. For $Re(\psi) \geq 0$, one has $\sigma_\psi(L) = \sigma_L(\xi)$, where $\xi = -\psi^2$. In *knotinfo* Web page [Ch-L], the parameter s satisfying $\xi = e^{\pi i s}$ is used. In particular, $\sigma_{6_2}(\xi) = -1$ for $Re(\xi) = \frac{3-\sqrt{5}}{4} = 1 - \cos(\pi/5) \approx 0.191$, and $s \approx 0.44$; compare [P-3].

⁹Khovanov homology and Heegaard-Floer homology allow construction of such signatures; e.g. Rasmussen invariant [Ras].

¹⁰Homology 4-ball glued along $K(1/n)$ to an NSPD homology sphere would produce a closed smooth 4-manifold with NSPD intersection form, contradicting Donaldson Theorem.

11, 1992) in which he demonstrated (using Kirby calculus) that $6_2(1/1)$ bounds a homology sphere with E_8 intersection form. Furthermore, according to [Co-G]: *Andreas Floer pointed out to us that if a homology sphere Σ bounds NSPD, then the Floer homology $I_*(\Sigma)$ is non-trivial, because Donaldson's Theorem will hold for X with boundary Σ as long as $I_*(\Sigma)$ is trivial.* $\square$

7. ACKNOWLEDGEMENTS

We would like to thank Selman Akbulut for providing us with the proof that $6_2(\frac{1}{1})$ bounds a homology sphere with E_8 intersection form.

8. AFTER TWENTY YEARS

The first version of this paper was written in 1990, almost 19 years before this arXiv version of the paper appears. Many results of the work (which has several abstracts published [P-1, P-2, T-3, T-4] and was available from the authors but was not published (and it was the time before arXiv), has been rediscovered and/or generalized. We should mention here the series of papers by A. Stoimenov [St-1, St-2, St-3, St-4, St-5], and papers by T. Nakamura [Nak], L. Rudolph [R-2], and M. Hirasawa [Hir]. Furthermore Ozawa proved that positive diagrams of a composite knots are visually composite [Oza] (see also [Cr]).

REFERENCES

- [Ca-G] A. J. Casson, C. McA. Gordon, On slice knots in dimension three, *Proceedings of Symposia in Pure Mathematics*, Volume 32, 1978, 39-53.
- [Ch-L] J. C. Cha, C. Livingston, KnotInfo: Table of Knot Invariants, <http://www.indiana.edu/knotinfo>.
- [Co-G] T. Cochran, E. Gompf, Applications of Donaldson's theorems to classical knot concordance, homology 3-spheres and property P , *Topology* 27 (1988), no. 4, 495-512.
- [Co-Li] T. Cochran, W. B. R. Lickorish, Unknotting information from 4-manifolds, *Trans. Amer. Math. Soc.*, 297(1), 1986, 125-142.
- [Con] J. H. Conway, An enumeration of knots and links, *Computational problems in abstract algebra* (ed. J. Leech), Pergamon Press, 1969, 329 - 358.
- [Cr-Mo] P. R. Cromwell, H. R. Morton, Positivity of knot polynomials on positive links, *J. Knot Theory Ramifications* 1(2), 1992, 203-206.
- [Cr] P. R. Cromwell, Positive braids are visually prime, *Proc. London Math. Soc.*, (3) 67, 1993, no. 2, 384-424.
- [Gi] C. A. Giller, A Family of links and the Conway calculus, *Trans. Amer. Math. Soc.*, 270(1), 1982, 75-109.
- [Gor] C. McA. Gordon, Some aspects of classical knot theory, *In: Knot theory*, L.N.M. 685, 1978, 1-60.
- [Hir] M. Hirasawa, Triviality and splittability of special almost alternating links via canonical Seifert surfaces, *Topology Appl.* 102, 2000, no. 1, 89-100.
- [Jo] V. F. R. Jones, Hecke algebra representations of braid groups and link polynomials, *Ann. of Math.* 126(2), 1987, 335-388.
- [Ka-1] L. H. Kauffman, *Formal knot theory*, Mathematical Notes 30, Princeton University Press, 1983 (the Dover edition 2006).
- [Ka-2] L. H. Kauffman, An invariant of regular isotopy, *Trans. Amer. Math. Soc.*, 318(2), 1990, 417-471.
- [Kir] R. Kirby, Problems in low-dimensional topology; Geometric Topology (Proceedings of the Georgia International Topology Conference, 1993), *Studies in Advanced Mathematics*, Volume 2 part 2., Ed. W. Kazez, AMS/IP, 1997, 35-473; (extended version of *Proc. Symp. Pure Math.*, 32, AMS, 1978, 35-56.
- [Ko] T. Kobayashi, Minimal genus Seifert surfaces for unknotting number 1 knots. *Kobe J. Math.* 6(1), 1989, 53-62.

- [KM] P. B. Kronheimer, T. S. Mrowka, Gauge theory for embedded surfaces. I, *Topology*, 32(4), 1993, 773–826.
- [L-T] W. B. R. Lickorish, M. B. Thistlethwaite, Some links with non-trivial polynomials and their crossing-numbers, *Comment. Math. Helv.*, 63, 1988, 527–539.
- [M-1] K. Murasugi, On a certain numerical invariant of link types. *Trans. Amer. Math. Soc.* 117, 1965, 387–422,
- [M-2] K. Murasugi, On the signature of links *Topology*, 9, 1970, 283–298.
- [M-3] K. Murasugi, On invariants of graphs with application to knot theory, *Trans. Amer. Math. Soc.*, 314, 1989, 1–49.
- [M-P-1] K. Murasugi, J. H. Przytycki, The Skein polynomial of a planar star product of two links, *Math. Proc. Cambridge Phil. Soc.*, 106, 1989, 273–276.
- [M-P-2] K. Murasugi, J. H. Przytycki, An index of a graph with applications to knot theory, *Memoirs of the American Math. Soc.*, Vol. 106, Number 508, November 1993, 101 pages.
- [Nak] Takuji Nakamura, Four-genus and unknotting number of positive knots and links, *Osaka J. Math.* 37, 2000, no. 2, 441–451.
- [Oza] Makoto Ozawa, Closed incompressible surfaces in the complements of positive knots, *Comment. Math. Helv.*, 77, 2002, 235–243.
- [P-1] J. H. Przytycki, Survey on recent invariants in classical knot theory, Warsaw University Preprints 6,8,9; 1986 (in English); a part of the book: *Knots: combinatorial approach to the knot theory*, Warsaw 1995 (in Polish);
e-print: <http://front.math.ucdavis.edu/0810.4191>
- [P-2] J. H. Przytycki, Positive knots have negative signature, *Bull. Ac. Pol. Math.* 37, 1989, 559–562. See also *Abstracts of AMS*, January 1988, vol. 9 No. 1, p. 147, where the conjecture about non-positivity of the signature of a link with a diagram with two negative crossings (with exception of a negative Hopf link, plus possible trivial components) is formulated. Corollary 1.10 proves and generalizes this conjecture.
- [P-3] J. H. Przytycki, Goeritz and Seifert matrices, Chapter IV of the book: **KNOTS: From combinatorics of knot diagrams to the combinatorial topology based on knots**, Cambridge University Press, accepted for publication, to appear 2011, pp. 600;
e-print: **arXiv Spring 2009; to appear**
- [P-Ta] J. H. Przytycki, K. Taniyama, Almost positive links have negative signature, *Abstracts of AMS*, June 1991, Issue 75, Vol. 12(3), p.327, *91T-57-69.
- [P-Tr-1] J. H. Przytycki, P. Traczyk, Conway algebras and skein equivalence of links, Preprint 1/86, Uniwersytet Warszawski, Instytut Matematyki, Warszawa 1986, 14 pages.
Partially published in [P-Tr-2].
- [P-Tr-2] J. H. Przytycki, P. Traczyk, Conway algebras and skein equivalence of links, *Proc. Amer. Math. Soc.*, 100(4), 1987, 744–748.
- [P-Tr-3] J. H. Przytycki, P. Traczyk, Supersignature and Milnor conjecture, unfinished manuscript.
- [Ras] J. Rasmussen, Khovanov homology and the slice genus,
<http://xxx.lanl.gov/abs/math.GT/0402131>
- [R-1] L. Rudolph, Nontrivial positive braids have positive signature, *Topology* 21(3), 1982, 325–327.
- [R-2] L. Rudolph, Positive links are strongly quasipositive, *Geometry and Topology Monographs*, 2: Proceedings of the Kirbyfest, 1999
- [S-T] M. Scharlemann, A. Thompson, Link genus and the Conway moves, *Comment. Math. Helv.*, 64(4), 1989, 527–535.
- [Sch] H. Schubert, Knoten mit zwei Brücken, *Math. Z.*, 65, 1956, 133–170.
- [St-1] A. Stoimenov, The signature of 2-almost positive knots. *J. Knot Theory Ramifications*, 9(6), 2000, 813–845.
- [St-2] A. Stoimenov, Gauss diagram sums on almost positive knots, *Compos. Math.* 140, 2004, no. 1, 228–254.
- [St-3] A. Stoimenov, On some restrictions to the values of the Jones polynomial, *Indiana Univ. Math. J.*, 54, 2005, no. 2, 557–574.
- [St-4] A. Stoimenov, Genus generators and the positivity of the signature. *Algebr. Geom. Topol.* 6 (2006), 2351–2393.
- [St-5] A. Stoimenov, Bennequin’s inequality and the positivity of the signature. *Trans. Amer. Math. Soc.*, 360(10), 2008, 5173–5199.

- [T-1] K. Taniyama, A partial order of knots, *Tokyo J. Math.* 12(1), 1989, 205-229.
- [T-2] K. Taniyama, A partial order of links, *Tokyo J. Math.* 12(2), 1989, 475-484.
- [T-3] K. Taniyama, Musubime no hanjunjo to sono ohyo (A partial order of knots and its applications). Proceedings of the 38th Topology Symposium, Kumamoto University, July 18-20, 1991, 81-96 (in Japanese).
- [T-4] K. Taniyama, Almost positive links have negative signature, Abstracts of talks presented at the Topology Section of the Annual Meeting of the Japanese Mathematical Society, April 1991, 10-11.
- [T-5] K. Taniyama, Links with two negative crossings, Abstracts of talks presented at the Topology Section of the Annual Meeting of the Japanese Mathematical Society, October 1991, 6-7.
- [Tr] P. Traczyk, Nontrivial negative links have positive signature. *Manuscripta Math.* 61 (1988), no. 3, 279-284.
- [Yo] Y. Yokota, Polynomial invariants of positive links, *Topology* 31(4), 1992, 805-811.

Kouki Taniyama
 e-mail: taniyama@waseda.jp
 Waseda University and GWU

Józef H. Przytycki
 e-mail: przytyck@gwu.edu
 George Washington University